

# FEDERAL REPUBLIC OF GERMANY

---



## Priority Certificate for the filing of a Patent Application

**File Reference:** 10 2004 014 280.7

**Filing date:** 22 March 2004

**Applicant/Proprietor:** Degussa AG, 40474 Düsseldorf/DE

**Title:** Process for preparing optically active amino acids  
using a whole-cell catalyst

**IPC:** C 12 P, C 12 N

The attached documents are a correct and accurate reproduction of the original submission for this application.


Munich, 25 February 2005  
**German Patent and Trademark Office**  
**The President**  
pp

[seal of the  
German  
Patent and  
Trademark Office]

[signature]  
Dzierzon

Process for preparing optically active amino acids  
using a whole-cell catalyst

The invention describes a process for preparing  
5 optically active L- $\alpha$ -amino acids. In particular the  
present invention describes a process for preparing  
compounds of the general formula (I)



(I),

10


in which R is alkyl, in particular a space-filling  
branched alkyl group which exhibits a tertiary C atom  
and which possesses 5-10 C atoms, for example tert-  
butyl, and substituted alkyl, or salts which are  
15 derived therefrom.

Optically active L- $\alpha$ -amino acids are used for preparing  
a number of valuable compounds. For example, these  
compounds function as intermediates in the production  
20 of pharmaceuticals. L-tert-Leucine, which can be found  
as a structural element in a number of pharmaceutical  
active compounds and is consequently required as an  
intermediate for synthesizing the corresponding  
pharmaceutical active compounds, is a particularly  
25 valuable representative of this product class.  
A. S. Bommarius et al., (J. Mol. Cat. B: Enzymatic  
1998, 5, 1-11) provides examples of uses of L-tert-  
leucine as a building block for pharmaceutical active  
compounds.

30

Using a leucine dehydrogenase and a formate  
dehydrogenase from *Candida boidinii* to enzymically  
reduce 2-ketocarboxylic acids while regenerating  
cofactor in situ constitutes an industrially

established method for preparing optically active L- $\alpha$ -amino acids. In particular, this route is suitable for preparing the non-proteinogenic amino acid L-tert-leucine, which is produced on the ton scale using this biocatalytic method. The method is described in detail in the literature (EP0692538; U. Kragl, D. Vasic-Racki, C. Wandrey, *Bioprocess Engineering* 1996, 14, 291-297; A. S. Bommarius, M. Schwarm, K. Drauz, *J. Mol. Cat. B: Enzymatic* 1998, 5, 1-11; G. Krix, A. S.: Bommarius, K. Kottenhahn, M. Schwarm, M.-R. Kula, *J. Biotechnol.* 1997, 53, 29-39, A. Liese, C. Wandrey, A. Liese, K. Seelbach, C. Wandrey, *Industrial Biotransformations*, Wiley-VCH, Weinheim, 2000, p. 125f. and A. S. Bommarius, K. Drauz, W. Hummel, M. -R. Kula, C. Wandrey, *Biocatalysis* 1994, 10, 37-47. In addition, a general review is provided in A. S. Bommarius in: *Enzyme Catalysis in Organic Synthesis* (Eds.: K. Drauz and H. Waldmann), Volume 2, 2nd edition, Wiley-VCH, Weinheim, 2003, chapter 15.3, p. 1047f.).



Scheme 1. Preparation of L-tert-leucine using isolated enzymes and added cofactor (taking as an example an  $\text{NAD}^+$ -dependent amino acid dehydrogenase and a formate

dehydrogenase for regenerating cofactor)

Typical quantities of NAD<sup>+</sup> cofactor which are used, and which have to be added, are described, for example, in  
5 EP0692538 and are in the range of from 0.0008 equivalents to 0.02 equivalents. In addition, G. Krix et al. (J. Biotechnol. 1997, 53, 29-39) describe the preparation of (S)-neopentylglycine in industrial batch sizes using an NAD<sup>+</sup> cofactor quantity of 0.003  
10 equivalents. Typical substrate concentrations in EP0692538 are 100-250 mM. A. Liese et al. (Industrial Biotransformations, Wiley-VCH, Weinheim, 2000, p. 125f.) describe the preparation of L-tert-leucine using a substrate concentration of 0.5 M and with a  
15 yield of 74%. G. Krix et al. (J. Biotechnol. 1997, 53, 29-39) also describe the performance of reductive aminations using isolated leucine dehydrogenase and formate dehydrogenase enzymes at substrate concentrations of from 0.5 to 1 M.

20

The high turnovers and outstanding enantioselectivities, which are > 99% ee and consequently help to meet the strict quality demands placed on pharmaceutical intermediates, are advantageous features of this  
25 method. It is also possible to operate at high substrate concentrations, something which is an important aspect particularly from the industrial point of view.

30 However, a disadvantage of the previous method is, in the first place, the requirement for isolated enzymes. These latter are used, in particular, in purified form, with this being accompanied by an increase in the share of the costs due to the biocatalyst. Because of the  
35 high enzyme costs resulting from this, it is necessary to recycle the enzymes many times in order to obtain a favorable process economy, in particular low enzyme costs. In addition to the long running times of these

recycling procedures, which are advantageously carried out continuously, the relatively small reaction volumes per batch which result from this are disadvantageous.

5 Another disadvantage is the requirement for cofactor which is added in the reaction. While these cofactors are added catalytically in orders of size of approx. 0.001 equivalents, they nevertheless represent, because of their high price, a considerable cost factor even at  
10 catalytic quantities.

A process in which the necessity of using isolated enzymes and of adding cofactor is dispensed with, or the addition of cofactor is kept to a minimum, and the  
15 synthesis nevertheless proceeds with a high turnover rate, high enantioselectivity and high volumetric productivity, would therefore be desirable. In this way, it would be possible to lower enzyme costs considerably and save on cofactor costs, and  
20 consequently increase the economy of the process.

Soda et al. describe the use of a whole-cell catalyst, comprising a leucine dehydrogenase and a bacterial formate dehydrogenase, in the reductive amination of,  
25 inter alia, branched-chain  $\alpha$ -ketocarboxylic acids such as L-tert-leucine (Appl. Environm. Microbiology 1997, 63, 4651-4656). This publication explicitly points out that the enzymes which are required in the reductive amination can be used in the form of a whole-cell  
30 catalyst, in particular E. coli, as live or resting cells, which comprises these enzymes. However, if preference were to be given to taking advantage of the intracellular pool of  $\text{NAD}^+$  in E.coli, for the purpose of avoiding having to add the  $\text{NAD}^+$ , the final  
35 concentration of product would then be restricted to about 0.3 M. This is not adequate for industrial applications.

The object of the present invention was therefore to specify another process for preparing L- $\alpha$ -amino acids which operates enzymically and which can be carried out advantageously on an industrial scale. The process  
5 should, in particular, be superior to the processes of the prior art with the abovedescribed aspects and should make it possible to produce the desired products advantageously from the point of view of process economics (in particular space-time yield).

10 These objects, and other objects which are not specified in more detail but which ensue from the prior art in an obvious manner, are achieved by a process having the features of the present claim 1. Claims 2 to  
15 9 are directed preferred embodiments of the present process.

Said object is achieved, in a manner which is extremely elegant and surprising but nonetheless advantageous for  
20 that, by, in a process for preparing enantiomerically enriched L- $\alpha$ -amino acids or their salts by reacting the corresponding 2-ketocarboxylic acid with an ammonium ion donor in the presence of a whole-cell catalyst which comprises a cloned gene encoding a cofactor-  
25 dependent amino acid dehydrogenase and a cloned gene encoding an enzyme which regenerates the cofactor, metering, at a total input of substrate per reaction volume of  $\geq 500$  mM, the addition of the substrate such that the stationary concentration of 2-ketocarboxylic  
30 acid is less than 500 mM and the external addition of cofactor, based on the total input of substrate, corresponds to  $< 0.0001$  equivalents.

Surprisingly, it is possible, for example by using the whole-cell catalyst while at the same time metering in  
35 the substrate, to dispense with any addition of the expensive cofactor or, by means of making a minimal external addition ( $< 0.0001$  equivalents), to keep its concentration in a low range, with this helping to save

on process input costs. By contrast, without this metering technology and when initially introducing substrate quantities per reaction volumes of > 500 mM, the reductive amination using the whole-cell catalyst only succeeds when relatively large quantities of the NAD<sup>+</sup> cofactor are added. In the absence of the latter, the concentration only proceeds unsatisfactorily (see comparative example "synthesis example 1", initial substrate quantity per reaction volumes 900 mM - final turnover 25%). It is consequently only by using the process according to the invention (see synthesis examples 2 to 4) that it is possible to be able to almost completely dispense with the external addition of the cofactor even when carrying out the synthesis with relatively high total turnover quantities per reaction volumes and consequently under conditions which make sense from the point of process economics.

In a preferred embodiment, the expensive cofactor is therefore only added in quantities which are such that a concentration of preferably < 0.00005 equivalents, extremely preferably < 0.00001 equivalents, based on the substrate, is maintained. Very particular preference is given to an embodiment in which no cofactor is added externally to the reaction mixture. In this case, therefore, no addition of the cofactors (e.g. NAD(H)) need take place at all, something which it was not possible to deduce in an obvious manner from the prior art.

Within the context of the reaction under consideration, the skilled person is free to choose the genes which encode a cofactor-dependent amino acid dehydrogenase and an enzyme which regenerates the cofactor, which genes are to be expressed by the whole-cell catalyst, as host organism. He will lean toward enzymes which are known from the prior art.


With regard to the amino acid dehydrogenase, suitable enzymes are, in particular, those which are selected from the group consisting of leucine dehydrogenases (US5854035) and phenylalanine dehydrogenases (US5416019). Amino acid dehydrogenases which have proved to be suitable are, in particular, the leucine dehydrogenases, with the leucine dehydrogenases from *Bacillus* strains, and, in this case, in particular, from *Bacillus sphaericus*, *Bacillus cereus* (Seq. ID No. 5) and *Bacillus stearothermophilus* being particularly suitable. Cofactor-regenerating enzymes which can be taken into consideration are those selected from the group consisting of formate dehydrogenases (EP1295937), malate dehydrogenases (PCT/EP/03/08631), lactate dehydrogenases and glucose dehydrogenases (the latter in A. Bommarius in: Enzyme Catalysis in Organic Synthesis (eds.: K. Drauz and H. Waldmann), Volume III, Wiley-VCH, Weinheim, 2002, chapter 15.3). The use of a formate dehydrogenase from *Candida boidinii* or mutants resulting therefrom (EP1295937; Seq. ID No. 7), while employing a formate-containing component as substrate, has proved to be very particularly preferred.

In this connection, a whole-cell catalyst which comprises a leucine dehydrogenase and a formate dehydrogenase from *Candida boidinii* or mutants derived therefrom is particularly suitable.

The substrate spectrum which is converted by the whole-cell catalyst differs depending on the amino acid dehydrogenase which is employed. While the leucine dehydrogenase is more suitable for linear and branched aliphatically substituted 2-ketocarboxylic acids, the phenylalanine dehydrogenase is preferably used for aromatic substituted substrates. With regard to the use of leucine dehydrogenase in the whole-cell catalyst, it is preferably possible to employ and convert substrates




of the general formula (II) possessing an aliphatic radical R



5

Substrates which possess bulky aliphatic radicals as R are particularly suitable. These R radicals are primarily those selected from the group consisting of 1-adamantyl, neopentyl and tert-butyl. For this reason, preference is given to a process in which use is made of 2-ketocarboxylic acids, or salts resulting therefrom, which yield amino acids of the general formula (I)



15

in which R is alkyl, in particular a space-filling branched alkyl group which exhibits a tertiary C atom and possesses 5-10 C atoms, for example tert-butyl, and substituted alkyls.

20

In principle, the skilled person is free to choose the manner in which he carries out the process according to the invention. In this connection, he will lean toward processes which are known from the prior art. These processes can be continuous or discontinuous. It is advantageous to meter the addition of the substrate in accordance with a fed batch process [see, for example, synthesis examples 2 and 4] or by continuously adding it [see, for example, synthesis example 3]. In both

25

30

process variants, the substrate is added such that the stationary concentration of substrate is less than 500 mM.

5 It has turned out to be advantageous to use the 2-keto-carboxylic acid employed as substrate at a maximum stationary concentration of less than 450 mM, and very particularly preferably of less than 400 mM, during the reaction.

10

In the fed batch process, the substrate is added in portions, after given units of time and preferably as a substrate solution. The number of the substrate portions which are added is preferably between 3 and 15, very preferably between 5 and 9. The concentration of the added substrate solution should preferably be set high enough to achieve a total input of substrate per reaction volume which is as high as possible. Synthesis examples 2 and 4 provide examples of this fed batch process variant. In the case of the continuous process variant, the substrate is added continuously over a given period of time, preferably at a constant metering rate, with the substrate preferably being added in the form of a substrate solution. Synthesis example 3 provides an example of this continuous process variant.

All known cells are suitable for use as the whole-cell catalyst which comprises an amino acid dehydrogenase and an enzyme which is capable of regenerating the cofactor. Microorganisms which may be mentioned in this regard are organisms such as yeasts, such as *Hansenula polymorpha*, *Pichia* sp., *Saccharomyces cerevisiae*, prokaryotes, such as *E.coli* and *Bacillus subtilis*, or eukaryotes, such as mammalian cells, insect cells or plant cells. The methods for cloning are well-known to the skilled person (Sambrook, J.; Fritsch, E. F. and Maniatis, T. (1989), Molecular cloning: a laboratory

manual, 2nd ed., Cold Spring Harbor Laboratory Press, New York). Preference is given to using *E.coli* strains for this purpose. Those which are very particularly preferred are: *E.coli* XL1 Blue, NM 522, JM101, JM109, JM105, RR1, DH5 $\alpha$ , TOP 10-, HB101, BL21 codon plus, BL21 (DE3) codon plus, BL21, BL21 (DE3), MM294. Plasmids which can preferably be used to clone the gene construct containing the nucleic acid according to the invention into the host organism are likewise known to the skilled person (see also PCT/EP03/07148; see below).

Suitable plasmids or vectors are, in principle, all the versions which are available to the skilled person for this purpose. These plasmids and vectors can be found, for example, in Studier and coworkers (Studier, W. F.; Rosenberg A. H.; Dunn J. J.; Dubendroff J. W.; (1990), Use of the T7 RNA polymerase to direct expression of cloned genes, *Methods Enzymol.* 185, 61-89) or the brochures provided by the companies Novagen, Promega, New England Biolabs, Clontech or Gibco BRL. Other preferred plasmids and vectors can be found in: Glover, D. M. (1985), *DNA cloning: a practical approach*, Vol. I-III, IRL Press Ltd., Oxford; Rodriguez, R. L. and Denhardt, D. T. (eds) (1988), *Vectors: a survey of molecular cloning vectors and their uses*, 179-204, Butterworth, Stoneham; Goeddel, D. V. (1990), *Systems for heterologous gene expression*, *Methods Enzymol.* 185, 3-7; Sambrook, J.; Fritsch, E. F. and Maniatis, T. (1989), *Molecular cloning: a laboratory manual*, 2nd ed., Cold Spring Harbor Laboratory Press, New York. Plasmids which can very preferably be used to clone the gene constructs containing the nucleic acid sequences under consideration into the host organism are, or are based on: pUC18/19 (Roche Biochemicals), pKK-177-3H (Roche Biochemicals), pBTac2 (Roche Biochemicals), pKK223-3 (Amersham Pharmacia Biotech), pKK-233-3 (Stratagene or pET (Novagen)).


In another embodiment of the process according to the invention, before it is used, the whole-cell catalyst is preferably pretreated such that the permeability of the cell membrane for the substrates and products is increased as compared with the intact system. In this connection, particular preference is given to a process in which the whole-cell catalyst is, for example, pretreated by being frozen and/or by being treated with toluene. The essential features of the process according to the invention are shown in scheme 2.

The substrates can be employed at an extraordinarily high concentration when using the present process, as has also been described in the prior art when using the individual enzymes. In the present case, it is advantageous to employ the 2-ketocarboxylic acid at a concentration of greater than 500 mM. It is also preferred to introduce the substrate into the reaction at concentrations of greater than 800 mM, preferably greater than 900 mM and very particularly preferably greater than 1000 mM. However, in the case of this embodiment, it is essential to add cofactor to the reaction mixture in order to achieve corresponding turnover rates.

If, however, it is wished, despite a high space-time yield being demanded, to use the whole-cell catalyst such that it does not become necessary to add the expensive cofactor externally, or only necessary to make an extremely small external addition of less than 0.0001 equivalents, the skilled person can then surprisingly achieve this by the metering, in accordance with the invention, of the substrate.

In the case of the present reaction, the procedure is preferably that the whole-cell catalyst and the ammonium ion donor are initially introduced in water. Any compound which is suitable to the skilled person for this purpose can be used as the ammonium ion donor.

In particular, these ammonium ion donors are compounds which are selected from the group consisting of typical ammonium salts. Very particular preference is given to using ammonium formate when a formate dehydrogenase is selected as the cofactor regeneration system. The reaction can be depicted very clearly by means of the following scheme 2.



Scheme 2. Principle of the reaction in the whole-cell catalyst process according to the invention (taking as an example an NAD<sup>+</sup>-dependent amino acid dehydrogenase and a formate dehydrogenase for regenerating cofactor)

If other dehydrogenases are used instead of the leucine dehydrogenase, the conditions under which the enzyme in question functions optimally can be found in the prior art. The reader is referred to US5416019 and Galkin et al. (Appl. Environ. Microbiol. 1997, 63, 4651) with regard to using a phenylalanine dehydrogenase.

With regard to the cofactor-regenerating enzymes and the conditions to be established, reference can be made to EP1295937 (formate dehydrogenase), PCT/EP/03/08631 (malate dehydrogenase) and Enzyme Catalysis in Organic Synthesis (Eds.: K. Drauz and H. Waldmann), Volume III, Wiley-VCH, Weinheim, 2002 (glucose dehydrogenase), and to the literature cited therein.

The reaction mixture is worked up using methods known to the skilled person. In the batch process, the biomass can be readily separated from the product by means of filtration or centrifugation. The amino acid  
5 which is obtained can then be isolated using customary methods (ion exchange chromatography, crystallization).

However, the present process can also be carried out continuously. For this, the reaction is carried out in  
10 what is termed an enzyme-membrane reactor in which high molecular weight substances, i.e. the biomass, are retained behind an ultrafiltration membrane and low molecular weight substances, such as amino acids which have been produced, are able to pass through the  
15 membrane. A procedure of this nature has already been described several times in the prior art (Wandrey et al. in year-book 1998, Verfahrenstechnik und Chemieingenieurwesen [Process technology and chemical engineering], VDI, p. 151ff; Kragl et al., Angew. Chem.  
20 1996, 6, 684).

The process, which is presented here, for preparing amino acids, which are, in particular, bulky, can very readily be established on a commercial scale on account  
25 of its advantages. The surprising fact that the addition, which is necessary in the case of the reaction under consideration, of a cofactor can be dispensed with in the process according to the invention, as well as the advantages arising from the  
30 fact that the whole-cell catalysts are easy to manage, constitute the non-obvious superiority of the present invention over the methods of the prior art.

Furthermore, it can be regarded as being surprising  
35 that the influence of undesirable metabolic/physiological functions is of no importance when using the whole-cell catalyst. Both aspects help, in an extraordinarily comprehensive manner, to lower the

process costs entailed in preparing the L- $\alpha$ -amino acids.

It is furthermore surprising that, despite permeabilization of the cell wall and the possibility, associated therewith, of the cofactor present in the cells escaping, a negative impairment of the reaction which might be expected, for example as a result of the turnover being decreased, is not observed.

Within the context of the invention, optically enriched (enantiomerically enriched, enantiomer enriched, enantiomerically pure) compounds are understood as meaning the presence of one optical antipode at > 50 mol% when mixed with the other.

The whole-cell catalyst is understood as meaning a microorganism which comprises cloned genes which encode enzymes which are at least able to catalyze two consecutive steps in the transformation of an organochemical compound. In this regard, and with regard to the general preparation methods (matching the enzyme expression with regard to the turnover rates), the reader is referred to EP1216304.

According to the invention, alkyl is understood as meaning a (C<sub>1</sub>-C<sub>18</sub>)-alkyl radical. This encompasses linear and arbitrarily branched radicals of this nature. It includes, in particular, methyl, ethyl, 1-propyl, 2-propyl, 1-n-butyl, 2-n-butyl, 1- or 2-isobutyl, 1- or 2-sec-butyl, tert-butyl, etc. The radicals can be substituted once or more than once by (C<sub>1</sub>-C<sub>8</sub>)-heteroalkyl radicals or radicals such as OH, SH, Hal and NH<sub>2</sub>. Heteroalkyl radicals are understood as meaning, in particular, an alkyl radical as described above which possesses from 1 to 8 C atoms and which contains heteroatoms, such as O, S or N in its chain or

which is bonded, by way of these heteroatoms, to the molecule under consideration.

5 External addition of cofactor means that this quantity of cofactor is added artificially to the reaction mixture. This quantity is to be seen as being in addition to the quantity of cofactor which is already inherently introduced into the reaction mixture by the whole-cell catalyst.

10

It goes without saying that the 2-ketocarboxylic acid which is used in the reaction is present in the reaction mixture in dissociated form. This form can be obtained either by using the ketocarboxylic acid and  
15 adjusting the pH correspondingly or by adding the salts of the ketocarboxylic acids. Both forms are included here analogously and in accordance with the invention.

20 The term total substrate concentration stands for the total input of substrate per reaction volume.



Figures:

Fig. 1 - pAM3.25 (Seq. ID No. 9):

5 Construction of pJOE4580.2

The plasmid pJOE4580.2 was formed from the published  
plasmid pJOE3075 (T. Stumpp, B. Wilms and  
J. Altenbuchner (2000) Biospektrum 1/2000: 33-36) by  
10 removing the male gene by cutting with the restriction  
endonucleases NdeI/HindIII and replacing it with two  
oligonucleotides which once again complemented the NdeI  
and HindIII cleavage sites and, in addition to this,  
carried an NheI, an AatII and a PstI cleavage site. A  
15 SmaI fragment from the plasmid pJOE773  
(J. Altenbuchner, P. Viell, I. Pelletier (1992)  
Positive selection vectors based on palindromic DNA  
sequences. Methods Enzymol 216: 457-466), which  
fragment carries the E.coli lacZalpha gene, was  
20 inserted into the NheI cleavage site after filling  
using Klenow polymerase and dNTPs. When harboring this  
plasmid, E. coli JM109 gives blue colonies on LB plates  
containing X-Gal and IPTG. This plasmid was named  
pJOE4580.2. The FDH sequence (Seq. ID No. 7) was cloned  
25 into this plasmid. The resulting plasmid was named  
pAM3.25.

Fig. 2 - pAM5.22

30 Construction of pJOE4580.2

The plasmid pJOE4580.2 was formed from the published  
plasmid pJOE3075 (T. Stumpp, B. Wilms and  
J. Altenbuchner (2000) Biospektrum 1/2000: 33-36) by  
35 removing the male gene by cutting with the restriction  
endonucleases NdeI/HindIII and replacing it with two  
oligonucleotides which once again complemented the NdeI  
and HindIII cleavage sites and, in addition to this,

carried an NheI, an AatII and a PstI cleavage site. A SmaI fragment from the plasmid pJOE773 (J. Altenbuchner, P. Viell, I. Pelletier (1992) Positive selection vectors based on palindromic DNA sequences. Methods Enzymol 216: 457-466), which fragment carries the E.coli lacZalpha gene, was inserted into the NheI cleavage site after filling using Klenow polymerase and dNTPs. When harboring this plasmid, E. coli JM109 gives blue colonies on LB plates containing X-Gal and IPTG. This plasmid was named pJOE4580.2. The LeuDH sequence (Seq. ID No. 5) was inserted into this plasmid. The new plasmid is named pAM5.22.

Fig. 3 - pAM8.21

Construction of pHWG640.12 (Seq. ID No. 11)

Plasmid pHWG640.12 has not previously been published and its construction is therefore described as follows. This plasmid pHWG640.12 is constructed proceeding from the published plasmid pAW229 in a manner which is readily reworkable. Plasmid pAW229 is a pACYC184 derivative which contains a rhamnose promoter. Proceeding from pAW229 (B. Wilms, A. Wiese, C. Syltatk, R. Mattes, J. Altenbuchner (2001) J. Biotechnol 86: 19-30), the hyuC gene was excised from the plasmid with NdeI/HindIII and replaced with a PCR fragment which was cut with the same restriction enzymes and which contains the E. coli K12 scfA (malic enzyme) gene. The resulting plasmid was designated pHWG640.12. The LeuDH sequence was inserted into this plasmid. The new plasmid is named pAM8.21.

Fig. 4 - pAM10.1 (Seq. ID No. 10)

The scfA gene (Seq. ID No. 11) was deleted from plasmid pAM8.21. The new plasmid is named pAM10.1.

Fig. 5

Biocatalyst with depiction of the course of the specific activity of leucine dehydrogenase (LeuDH) and formate dehydrogenase (FDH), and of the optical density, in dependence on the induction time; for a detailed description of the fermentation conditions, see experimental section.

#### Experimental examples

10

Preparing the whole-cell catalyst

#### Gene amplification and cloning

15 In order to clone the formate dehydrogenase (FDH, *fdh3* from *Candida boidinii*, mutant with lower sensitivity to oxidation) and leucine dehydrogenase (*Bacillus cereus* LeuDH) for the whole-cell catalysis of the conversion of trimethylpyruvate into tert-leucine with  
20 regeneration of cofactor, the genes for the two enzymes were first of all amplified by PCR from chromosomal DNA obtained from the abovementioned strains. The oligonucleotides employed are listed in Table 1 while the composition of the PCR mixtures is given in Table 2  
25 and the PCR program is given in Table 3.

Table 1: Oligonucleotides for amplifying the FDH and LeuDH genes

Oligo-nucleotide	5'-3' sequence		Seq. ID No.
s3713	AAA AAA <u>CTT AAG</u> AAG GAG ATA TAC ATA TGA CAT TAG AAA TCT TCG AA	LeuDH forward	1
s3714	AAA AAA <u>CTG CAG</u> TTA GCG ACG GCT AAT AAT AT	LeuDH reverse	2
s3723	AAA AAA <u>CAT ATG</u> AAG ATT GTC TTA GTT CTT	FDH forward	3
s3716	AAA AAA <u>GAC GTC</u> TTA TTT CTT ATC GTG TTT ACC	FDH reverse	4

The oligonucleotides were used to append cleavage sites for restriction endonucleases to the genes. These are

5 BfrI in the case of s3713, PstI in the case of s3714, NdeI in the case of s3723 and AatII in the case of s3716 (see underlined regions).

10 Table 2: PCR mixtures, polymerase, buffer and MgCl<sub>2</sub> originate from the company Biomaster; the plasmid DNA starting concentration was 50 µg/ml

Component	For FDH	Mixture for FDH	for LeuDH	Mixture for LeuDH
Plasmid DNA from strain FDH-C235/C262A		2 µl	pLeu2 plasmid DNA	2 µl
10× buffer		10 µl		10 µl
50 mM MgCl <sub>2</sub>		3 µl		3 µl
100% DMSO		10 µl		10 µl
10 mM dNTPs		2 µl		2 µl
33 mM oligo 1	s3723	1 µl	s3713	1 µl
33 mM oligo 2	s3716	1 µl	s3714	1 µl
Taq polymerase		1 µl		1 µl
Deionized H <sub>2</sub> O		70 µl		70 µl

Table 3: PCR program: steps 2 to 4 were repeated 30 times

Step	T, t for FDH amplification	T, t for LeuDH amplification
1. Denaturation of the DNA	94°C, 5 min	94°C, 5 min
2. Oligo annealing	50°C, 1 min	51°C, 1 min
3. DNA elongation	72°C, 1: 30 min	72°C, 1:30 min
4. Denaturation of the dsDNA	92°C, 1 min	92°C, 1 min
5. DNA elongation	72°C, 7 min	72°C, 7 min

5 After the gene amplification, the PCR fragments were purified using the "DNA PCR and gel band purification kit" supplied by the company GFX and ligated into the L-rhamnose-inducible vectors pJOE4580.2 (pBR322 derivative; Fig. 1) and, respectively, pHWG640.12  
10 (pACYC184 derivative; Fig. 3; Seq. ID No. 11) using the restriction endonucleases mentioned below.

In general, restriction mixtures were prepared using approx. 50 µg of DNA/ml in the 10 µl standard mixture.  
15 1 µl of the first enzyme, and 1 µl of the 10x concentrated enzyme buffer, were also added. The mixtures were adjusted to the final volume using deionized H2O. The DNA to be inserted was incubated with the restriction enzymes separately from the  
20 plasmid DNA. After the restriction with the first enzyme, there then followed a precipitation step in which the DNA was precipitated with isopropanol and washed with ethanol and then dried and taken up in 8 µl of TE 10.01. In each case 1 µl of the second enzyme and  
25 1 µl of the second 10x enzyme buffer were added to these mixtures, which were incubated once again at 37°C for 1.5 h. The vector pAM10.1 was prepared from pAM8.21, this was also followed by a treatment with

Klenow polymerase. The DNA was then separated into its fragments using a 1% agarose gel (Seakem agarose containing 0.4  $\mu$ g of ethidium bromide/ml) and the correct bands were excised with a scalpel for further use. The DNA was eluted, in accordance with the instructions, from the small gel blocks using the "EASY PURE gel purification kit" supplied by the company Biozym and taken up in 15  $\mu$ l TE 10.01.

For the ligation of vector and insert, the mixtures were selected such that the insert DNA was present at approximately twice the concentration of the target vector. In this case, too, the DNA concentration was approx. 50  $\mu$ g/ml. The final volume of the ligation mixtures was 10  $\mu$ l, with the mixtures also containing 1  $\mu$ l of ligase and 1  $\mu$ l of 10 $\times$  concentrated ligase buffer (both from ROCHE) in addition to the vector/insert mixture. The incubation took place overnight at 4°C. The ligation mixtures were transformed into E.coli K12 JM109, with this bacterium then being selected on LB agar containing antibiotics (100  $\mu$ g of ampicillin/ml (pAM3.25 [Seq. ID No. 9], pAM5.22) or 25  $\mu$ g chloramphenicol/ml (pAM8.21, pAM10.1 [Seq. ID No. 10])), and clones were checked for the expected plasmid after the plasmids had been isolated.

Since LeuDH (Seq. ID No. 6) was initially to be coupled to malic enzyme (Seq. ID No. 12), the LeuDH gene was first of all inserted into pJOE4625.1, which already contained the gene for malic enzyme (sfca) (Fig. 2). The LeuDH gene was then inserted into pHGW640.12 (Fig. 3), a pACYC184 derivative which also contained a rhamnose promoter and an sfca gene, which latter was then deleted. The subcloning of the LeuDH gene from plasmid pAM5.22 (Fig. 2) into the target plasmid pAM10.1 (Fig. 4) was necessary in order to construct a two-plasmid system which requires two resistance markers for selection.

Table 4: Cloning results

Gene/vector	Cloned into plasmid	Restriction with	New designation	Fig.
FDH PCR fragment	pJOE4580.2	NdeI, AatII	pAM3.25	1
LeuDh PCR fragment	pJOE4625.1	BfrI, PstI	pAM5.22	2
LeuDh from pAM5.22	pHWG640.12	BfrI, BamHI	pAM8.21	3
pAM8.21	Without sfcA gene	MunI, PstI	pAM10.1	4

## Fermenting the whole-cell catalyst

5 After HPLC analysis had shown that the FDH/LeuDh combination (*E.coli* JM109/pAM3.25/pAM10.1) achieved better results in converting trimethylpyruvate into tert-leucine than a comparative model system (malic enzyme/LeuDh on pAM5.22) in miniature-scale (1 ml) experiments in a thermoshaker, plasmids pAM3.25 and pAM10.1 were transformed into *E.coli* BW3110 since this strain is more suitable for fermentations. The intention was to use high cell density fermentation to prepare a sufficiently large biomass for all the following investigations using the model system. The fermentation was carried out without any antibiotic, with the preliminary cultures having been grown in the presence of antibiotic, at 30°C in a 30 l fermenter containing a final volume of 8 l. For this, the cells were initially grown at 30°C as a batch culture up to an OD600 = 50 and until the glucose had been completely consumed (approx. 22h). Gene expression was then induced by adding rhamnose, which had been sterilized by filtration, to a final concentration of 0.2%, while fed batch culture was started by automatically adding nutrient solution and minerals (feed I and feed II). Samples, whose OD and enzyme activities were

determined, using the respective activity tests in the latter case, were taken every two hours from the induction onward. The course of the OD, and of the activities, until fermentation was terminated are plotted against the time in Figure 5.

The fermentation was terminated 22h after the rhamnose induction since, despite increasing cell density, the activity of the FDH had stagnated and the cause of this was presumably plasmid loss or a reaction medium which was too acidic. The latter became apparent in the whole-cell reactions, in which the pH fell markedly ( $\Delta\text{pH}_{\text{max}} = 0.8$ ), as compared with a previously pH-regulated solution, when the moist biomass was added. The activities of the two enzymes reached 0.565 U/mg of total protein in the case of the LeuD<sub>H</sub> and 0.123 U/mg of total protein in the case of the FDH. The volume activities, based on the fermentation medium, were 32.77 U/ml for the LeuD<sub>H</sub> and 7.14 U/ml for the FDH. After the medium had been removed in a separator, the cell yield was 1.4 kg of moist biomass. The cells were stored temporarily at -20°C until being used as whole-cell catalyst.

## 25 Fermentation media

Preliminary culture: 2 x 200 ml

Preliminary culture medium:  $\text{cNa}_2\text{SO}_4 \times 10\text{H}_2\text{O} = 2 \text{ g/l}$

$\text{c}(\text{NH}_4)_2\text{SO}_4 = 2.675 \text{ g/l}$

$\text{cNH}_4\text{Cl} = 0.5 \text{ g/l}$

$\text{cK}_2\text{HPO}_4 = 14.625 \text{ g/l}$

$\text{cNaH}_2\text{PO}_4 \times 2\text{H}_2\text{O} = 3.6 \text{ g/l}$



autoclave in 90% by vol.  
H<sub>2</sub>O

cglucose = 10 g/l, final  
concentration

(stock solution in H<sub>2</sub>O)

autoclave separately

5

1M MgSO<sub>4</sub> solution, 2 ml/l

TES, 3 ml/l

Thiamine stock solution (10 g/l in H<sub>2</sub>O), 1 ml/l

- 10 Batch culture: Add inoculum (380 ml in which Cx = 12 g/l) containing glucose, MgSO<sub>4</sub>, TES and thiamine in an inoculation flask to the autoclaved batch medium

Batch medium (quantity taken for 8 l):

Na <sub>2</sub> SO <sub>4</sub> × 10H <sub>2</sub> O	16 g
(NH <sub>4</sub> ) <sub>2</sub> SO <sub>4</sub>	21.4 g
NH <sub>4</sub> Cl	4 g
K <sub>2</sub> HPO <sub>4</sub>	117 g
NaH <sub>2</sub> PO <sub>4</sub> × 2H <sub>2</sub> O	28.8 g
(NH <sub>4</sub> ) 2H-citrate	8 g

dissolve in 7.5 l of H<sub>2</sub>O and sterilize in a  
30 l fermenter

Glucose monohydrate	220 g
---------------------	-------

dissolve in 500 ml of H<sub>2</sub>O and autoclave  
(25 g/l)

1M MgSO <sub>4</sub> solution	16 ml
-------------------------------	-------

TES	24 ml
-----	-------

Thiamine solution (10 g/l)	8 ml
----------------------------	------

(sterilize the thiamine by filtration,  
autoclave the remainder)

pH 7.2, using H<sub>3</sub>PO<sub>4</sub> and NH<sub>3</sub>

## Fed batch feed:

I.    Glucose monohydrate            2750 g in 3.5 l of H<sub>2</sub>O  
         autoclave  
5       MgSO<sub>4</sub> × 7H<sub>2</sub>O                98.5 g in 0.15 l of H<sub>2</sub>O  
         autoclave  
         TES solution                0.5 l  
         autoclave  
         Thiamine                    2.5 g in 0.5 l of H<sub>2</sub>O  
10       sterilize by filtration  
         then combine in a feed flask

II.    (NH<sub>4</sub>)<sub>2</sub>HPO<sub>4</sub>                    396 g in 1 l of H<sub>2</sub>O, pH 7  
         autoclave

15    Feeds I and II are added using two separate pumps

pH:     7.2 (titrated with H<sub>3</sub>PO<sub>4</sub> and NH<sub>3</sub>)

20    pO<sub>2</sub>:    approx. 50 kPa (regulated by the rotational  
         speed of the agitator)

Trace element  
solution (TES):

	CaCl <sub>2</sub> × 2H <sub>2</sub> O	0.5 g
25	ZnSO <sub>4</sub> × 7H <sub>2</sub> O	0.18 g
	MnSO <sub>4</sub> × H <sub>2</sub> O	0.1 g
30	Di-Na-EDTA	20.1 g
	FeCl <sub>3</sub> × 6H <sub>2</sub> O	16.7 g
	CuSO <sub>4</sub> × 5H <sub>2</sub> O	0.16 g
35	CoCl <sub>2</sub> × 6H <sub>2</sub> O	0.18 g
	H <sub>2</sub> O to 1 l	

Preparing L-tert-leucine using a whole-cell catalyst at 900 mM without metering (comparative example = synthesis example 1)

5 50 ml of an 0.9 M solution of trimethylpyruvic acid (pH 7.0, adjusted with 32% ammonia), which also contains 1 mM magnesium chloride and 1% (v/v) toluene, are added to 5.85 g of the biocatalyst (*E.coli* JM105 (pAM 3.25\_10.1) biomass) and 7.95 g of ammonium formate  
10 (2.8 mol equivalents). The pH is adjusted to pH 7.0 at the beginning of the reaction and not regulated any further after that, resulting in the pH rising during the reaction. The reaction temperature is 30°C. After a reaction time of 8 h, a conversion of 24.6% is  
15 measured, with it not being possible to increase this conversion any further even after an additional 15 h of stirring.

Preparing L-tert-leucine using a whole-cell catalyst at  
20 approx. 0.9 M and employing fed batch metering (synthesis example 2)

23.84 g of ammonium formate (corresponding to 2.8 equivalents based on the total substrate quantity  
25 employed) and 17.55 g of the biocatalyst (*E.coli* JM105 (pAM 3.25\_10.1) biomass) are initially weighed into a 250 l three-neck flask, after which 28.50 ml of deionized water and 150 µl of a 1M solution of magnesium chloride (corresponding to a 1 mM  
30 concentration based on the final volume) are added. When the reaction temperature of 30°C has been reached, the reaction is started by adding 7.50 ml of a 1.8 M solution of trimethylpyruvic acid (pH 7.0, adjusted with 32% ammonia). The pH is then adjusted to 7.0 by  
35 adding 32% ammonia. After that, in each case 7.50 ml of a 1.8 M solution of trimethylpyruvic acid (pH 7.0, adjusted with 32% ammonia) are firstly metered in twice after which different volumes of a 1.8 M solution of

trimethylpyruvic acid (pH 7.0, adjusted with 32% ammonia) are metered in five times, with all the additions taking place at defined time intervals. The time intervals, and the quantities which are in each case metered in, are given in the following metering table. The final volume is 150 ml and the total concentration of added substrate is 0.86 M, corresponding to a volumetric quantity of trimethylpyruvic acid of 112.5 g/l. A complete conversion (> 98% in accordance with HPLC) is observed after a reaction time of 24 h.

Metering table	Substrate solution	Substrate solution
Time (h)	ml (1.8 M)	ml (0.9 M)
0	7.5	0
0.5	7.5	0
1	7.5	0
2.5	0	15
4	0	17.5
5.5	0	20
6.5	0	22.5
7	0	24
Total volume of metered-in substrate solution	22.5	99

Preparing L-tert-leucine using a whole-cell catalyst at 1 M and employing continuous metering (synthesis example 3)

26.48 g of ammonium formate (corresponding to 2.8 equivalents based on the total quantity of substrate employed), 150  $\mu$ l of a 1 M solution of magnesium chloride (corresponding to a 1 mM concentration based on the final volume) and 19.49 g of the biocatalyst (E.coli JM105 (pAM3.25\_10.1) biomass) are initially weighed into a 250 l three-neck flask, after which

30 ml of deionized water are added. The pH is then adjusted to 7.0 by adding 32% ammonia. After the reaction temperature of 30°C has been reached, a total of 120 ml of a 1.25 M solution of trimethylpyruvic acid (pH 7.0, adjusted with 32% ammonia) are added continuously at a flow rate of 0.2 ml/min over a period of 10 hours. The final volume is 150 ml and the total concentration of substrate employed is 1.0 M, corresponding to a volumetric quantity of trimethylpyruvic acid of 130.1 g/l. A conversion of 96% (in accordance with HPLC) is observed after a reaction time of 27 h.

Preparing L-tert-leucine using a whole-cell catalyst at 700 mM and employing fed batch metering

2.55 g of sodium formate (corresponds to 2.5 mol/l based on final volume) are initially added to a conically shaped 100 ml reaction flask belonging to a STAT Titrino 718, after which 15 µl of a 1 M solution of  $MgCl_2$  (corresponds to a final concentration of 1 mM) and 4.5 ml of a 1 M solution of TMP (pH 7, adjusted with 25% ammonia), and also 1.5% by vol. of toluene (based on the final volume), are added. The volume is made up to 15 ml with deionized  $H_2O$ . The reaction temperature of 30°C is kept stable, and controlled, by a closed-loop water circuit. 1 g of the biocatalyst moist biomass is resuspended in the substrate mixture and the pH is adjusted to 6.9 to 7 with 25% ammonia.

After pH 7.5 has been reached, 4.5 ml of the 1 M TMP solution (pH 7) are added repeatedly. In this connection, the pH falls by approx.  $\Delta pH = 0.3$ . As soon as pH 7.5 is reached, 4.5 ml of 1 M TMP solution are added once again. The addition of said volume of TMP is repeated 10x until the pH does not fall any further when TMP is added. In addition, 4 ml of a 4 M solution of sodium formate (corresponds, without taking any

reaction into consideration, to a concentration of 973 mM in the medium) are added in connection with the eighth addition of TMP. The final volume is 64 ml, with a volumetric final concentration (without taking the  
5 reaction into consideration) of trimethylpyruvic acid of 774 mM (100.6 g/l). Sodium formate is present in solution at a final concentration of 836 mM. HPLC showed that 92% of the trimethylpyruvic acid had been converted after only 6h.


10

The concentrations of the two substrates at the different addition points are listed in Table 5 below.

Time [t in min]	Concentration of trimethyl- pyruvic acid [mM]	Concentration of sodium formate [mM]	Second addition of sodium formate
0	300	2500	
45	461.54	1923.08	
60	562.5	1562.5	
75	631.58	1315.79	
90	681.82	1136.36	
105	720	1000	
120	750	892.86	
135	774.19	806.45	
150	736.36	972.73	x
180	756.30	899.16	
210	773.44	835.94	

## Patent claims:

1. A process for preparing enantiomerically enriched L- $\alpha$ -amino acids or their salts by reacting the corresponding 2-ketocarboxylic acid with an ammonium ion donor in the presence of a whole-cell catalyst which comprises a cloned gene encoding a cofactor-dependent amino acid dehydrogenase and a cloned gene encoding an enzyme which regenerates the cofactor, at a total input of substrate per reaction volume of  $\geq 500$  mM, with the addition of the substrate being metered such that the stationary concentration of 2-ketocarboxylic acid is less than 500 mM and the external addition of cofactor, based on the total input of substrate, corresponds to  $< 0.0001$  equivalents.
2. The process as claimed in claim 1, characterized in that no cofactor is added to the reaction mixture.
3. The process as claimed in claim 1 and/or 2, characterized in that use is made of 2-ketocarboxylic acids which yield amino acids of the general formula (I)




in which R is alkyl, in particular a space-filling branched alkyl group which exhibits a tertiary C atom and possesses 5-10 C atoms, for example tert-butyl, and substituted alkyls.

4. The process as claimed in one or more of the preceding claims,  
characterized in that  
the substrate is metered in in accordance with a  
5 fed batch process.
5. The process as claimed in one or more of the preceding claims,  
characterized in that  
10 the 2-ketocarboxylic acid is kept at a maximum stationary concentration of less than 450 mM, very preferably of less than 400 mM.
6. The process as claimed in one or more of the preceding claims,  
15 characterized in that  
before it is used, the whole-cell catalyst is pretreated such that the permeability of the cell membrane for the substrate and products is  
20 increased as compared with the intact system.



## Abstract:

The present invention relates to a process for preparing, in particular, enantiomerically enriched  
5 L- $\alpha$ -amino acids, in particular those of the general formula (I).



(I)

- 10 In this connection, the process according to the invention uses 2-ketocarboxylic acids which are converted into the desired products using a whole-cell catalyst which comprises an amino acid dehydrogenase and a cofactor-regenerating enzyme.

Fig. 1




Fig. 2




Fig. 3




Fig. 4





Fig. 5



## SEQUENCE LISTING

<110> Degussa AG

5 <120> Process for preparing optically active amino acids using a whole-cell catalyst

<130> 040055 AM

10 <160> 12

<170> PatentIn version 3.1

15 <210> 1  
<211> 47  
<212> DNA  
<213> Artificial

20 <220>  
<223> Primer

<400> 1  
aaaaaaactta agaaggagat atacatatga cattagaaat cttcgaa 47

25 <210> 2  
<211> 32  
<212> DNA  
<213> Artificial

30 <220>  
<223> Primer

<400> 2

35 aaaaaactgc agttagcgac ggctaataat at 32

<210> 3  
<211> 30  
40 <212> DNA  
<213> Artificial

<220>  
<223> Primer

45 <400> 3  
aaaaaacata tgaagattgt cttagttctt 30

50 <210> 4  
<211> 32  
<212> DNA  
<213> Artificial

55 <220>  
<223> Primer

<400> 4  
aaaaaagacg tcttatttct tatcgtgttt ac 32

```

<210> 5
<211> 1120
5 <212> DNA
   <213> Bacillus cereus

<220>
<221> CDS
10 <222> (20)..(1120)
   <223>

<400> 5
15 ttaagaagga gatatacat atg aca tta gaa atc ttc gaa tac tta gaa aaa 52
   Met Thr Leu Glu Ile Phe Glu Tyr Leu Glu Lys
   1 5 10

   tat gat tat gag caa gta gta ttt tgt caa gat aaa gaa tct ggt tta 100
   Tyr Asp Tyr Glu Gln Val Val Phe Cys Gln Asp Lys Glu Ser Gly Leu
   15 20 25

   aaa gca att att gca att cat gat aca aca ctt gga ccg gct ctt ggt 148
   Lys Ala Ile Ile Ala Ile His Asp Thr Thr Leu Gly Pro Ala Leu Gly
   30 35 40

25 gga aca aga atg tgg aca tat gat tct gaa gaa gcg gcg att gaa gat 196
   Gly Thr Arg Met Trp Thr Tyr Asp Ser Glu Glu Ala Ala Ile Glu Asp
   45 50 55

30 gca ttg cgt ctt gca aaa ggg atg aca tac aaa aac gca gca gct ggt 244
   Ala Leu Arg Leu Ala Lys Gly Met Thr Tyr Lys Asn Ala Ala Ala Gly
   60 65 70 75

35 tta aac tta ggt ggt gcg aaa aca gta att atc ggt gat cct cgt aaa 292
   Leu Asn Leu Gly Gly Ala Lys Thr Val Ile Ile Gly Asp Pro Arg Lys
   80 85 90

   gat aag agc gaa gca atg ttc cgt gca cta gga cgt tat atc caa gga 340
   Asp Lys Ser Glu Ala Met Phe Arg Ala Leu Gly Arg Tyr Ile Gln Gly
   95 100 105

40 cta aac gga cgt tac att aca gct gaa gat gtt ggt aca aca gta gat 388
   Leu Asn Gly Arg Tyr Ile Thr Ala Glu Asp Val Gly Thr Thr Val Asp
   110 115 120

45 gat atg gat att atc cat gaa gaa act gac ttt gta aca ggt atc tca 436
   Asp Met Asp Ile Ile His Glu Glu Thr Asp Phe Val Thr Gly Ile Ser
   125 130 135

50 cca tca ttc ggt tct tct ggt aac cca tct ccg gta act gca tac ggt 484
   Pro Ser Phe Gly Ser Ser Gly Asn Pro Ser Pro Val Thr Ala Tyr Gly
   140 145 150 155

55 gtt tac cgt ggt atg aaa gca gct gca aaa gaa gct ttc ggt act gac 532
   Val Tyr Arg Gly Met Lys Ala Ala Ala Lys Glu Ala Phe Gly Thr Asp
   160 165 170

aat tta gaa gga aaa gta att gct gtt caa ggc gtt ggt aac gta gca 580
Asn Leu Glu Gly Lys Val Ile Ala Val Gln Gly Val Gly Asn Val Ala

```



	175	180	185	
5	tat cac cta tgc aaa cat tta cac gct gaa gga gca aaa tta att gtt Tyr His Leu Cys Lys His Leu His Ala Glu Gly Ala Lys Leu Ile Val 190 195 200			628
10	aca gat att aat aaa gaa gct gta caa cgt gct gta gaa gaa ttc ggt Thr Asp Ile Asn Lys Glu Ala Val Gln Arg Ala Val Glu Glu Phe Gly 205 210 215			676
15	gca tca gca gtt gaa cca aat gaa att tac ggt gtt gaa tgc gat att Ala Ser Ala Val Glu Pro Asn Glu Ile Tyr Gly Val Glu Cys Asp Ile 220 225 230 235			724
20	tac gca cca tgt gca cta ggc gca aca gtt aat gat gaa act att cca Tyr Ala Pro Cys Ala Leu Gly Ala Thr Val Asn Asp Glu Thr Ile Pro 240 245 250			772
25	caa ctt aaa gca aaa gta atc gca ggt tct gcg aat aac caa tta aaa Gln Leu Lys Ala Lys Val Ile Ala Gly Ser Ala Asn Asn Gln Leu Lys 255 260 265			820
30	gaa gat cgt cat ggt gac atc att cat gaa atg ggt att gta tac gca Glu Asp Arg His Gly Asp Ile Ile His Glu Met Gly Ile Val Tyr Ala 270 275 280			868
35	cca gat tat gta att aat gca ggt ggc gta att aac gta gca gac gaa Pro Asp Tyr Val Ile Asn Ala Gly Gly Val Ile Asn Val Ala Asp Glu 285 290 295			916
40	tta tat gga tac aat aga gaa cgt gca cta aaa cgt gtt gag tct att Leu Tyr Gly Tyr Asn Arg Glu Arg Ala Leu Lys Arg Val Glu Ser Ile 300 305 310 315			964
45	tat gac acg att gca aaa gta atc gaa att tca aaa cgc gat ggc ata Tyr Asp Thr Ile Ala Lys Val Ile Glu Ile Ser Lys Arg Asp Gly Ile 320 325 330			1012
50	gca act tat gta gcg gca gat cgt cta gct gaa gag cgc att gca agc Ala Thr Tyr Val Ala Ala Asp Arg Leu Ala Glu Glu Arg Ile Ala Ser 335 340 345			1060
55	ttg aag aat tct cgt agc act tac tta cgc aac ggt cac gat att att Leu Lys Asn Ser Arg Ser Thr Tyr Leu Arg Asn Gly His Asp Ile Ile 350 355 360			1108
	agc cgt cgc taa Ser Arg Arg 365			1120
	<210> 6 <211> 366 <212> PRT <213> Bacillus cereus  <400> 6			

Met Thr Leu Glu Ile Phe Glu Tyr Leu Glu Lys Tyr Asp Tyr Glu Gln

```

1          5          10          15
5  Val Val Phe Cys Gln Asp Lys Glu Ser Gly Leu Lys Ala Ile Ile Ala
    20                      25                      30
10 Ile His Asp Thr Thr Leu Gly Pro Ala Leu Gly Gly Thr Arg Met Trp
    35                      40                      45
15 Thr Tyr Asp Ser Glu Glu Ala Ala Ile Glu Asp Ala Leu Arg Leu Ala
    50                      55                      60
20 Lys Gly Met Thr Tyr Lys Asn Ala Ala Ala Gly Leu Asn Leu Gly Gly
    65                      70                      75                      80
25 Ala Lys Thr Val Ile Ile Gly Asp Pro Arg Lys Asp Lys Ser Glu Ala
    85                      90                      95
30 Met Phe Arg Ala Leu Gly Arg Tyr Ile Gln Gly Leu Asn Gly Arg Tyr
    100                     105                     110
35 Ile Thr Ala Glu Asp Val Gly Thr Thr Val Asp Asp Met Asp Ile Ile
    115                     120                     125
40 His Glu Glu Thr Asp Phe Val Thr Gly Ile Ser Pro Ser Phe Gly Ser
    130                     135                     140
45 Ser Gly Asn Pro Ser Pro Val Thr Ala Tyr Gly Val Tyr Arg Gly Met
    145                     150                     155                     160
50 Lys Ala Ala Ala Lys Glu Ala Phe Gly Thr Asp Asn Leu Glu Gly Lys
    165                     170                     175
55 Val Ile Ala Val Gln Gly Val Gly Asn Val Ala Tyr His Leu Cys Lys
    180                     185                     190
60 His Leu His Ala Glu Gly Ala Lys Leu Ile Val Thr Asp Ile Asn Lys
    195                     200                     205
65 Glu Ala Val Gln Arg Ala Val Glu Glu Phe Gly Ala Ser Ala Val Glu
    210                     215                     220
70 Pro Asn Glu Ile Tyr Gly Val Glu Cys Asp Ile Tyr Ala Pro Cys Ala
    225                     230                     235                     240

```

	Leu Gly Ala Thr Val Asn Asp Glu Thr Ile Pro Gln Leu Lys Ala Lys	245	250	255
5	Val Ile Ala Gly Ser Ala Asn Asn Gln Leu Lys Glu Asp Arg His Gly	260	265	270
10	Asp Ile Ile His Glu Met Gly Ile Val Tyr Ala Pro Asp Tyr Val Ile	275	280	285
15	Asn Ala Gly Gly Val Ile Asn Val Ala Asp Glu Leu Tyr Gly Tyr Asn	290	295	300
20	Arg Glu Arg Ala Leu Lys Arg Val Glu Ser Ile Tyr Asp Thr Ile Ala	305	310	315
25	Lys Val Ile Glu Ile Ser Lys Arg Asp Gly Ile Ala Thr Tyr Val Ala	325	330	335
30	Ala Asp Arg Leu Ala Glu Glu Arg Ile Ala Ser Leu Lys Asn Ser Arg	340	345	350
35	Ser Thr Tyr Leu Arg Asn Gly His Asp Ile Ile Ser Arg Arg	355	360	365
	<210> 7			
	<211> 1095			
	<212> DNA			
	<213> Candida boidinii			
	<220>			
	<221> CDS			
	<222> (1)..(1095)			
	<223>			
	<400> 7			
45	atg aag att gtc tta gtt ctt tat gat gct ggt aag cac gct gct gat			48
	Met Lys Ile Val Leu Val Leu Tyr Asp Ala Gly Lys His Ala Ala Asp			
	1 5 10 15			
50	gaa gaa aaa tta tat ggt tct act gaa aat aaa tta ggt att gct aat			96
	Glu Glu Lys Leu Tyr Gly Ser Thr Glu Asn Lys Leu Gly Ile Ala Asn			
	20 25 30			
55	tgg tta aaa gat caa ggt cat gaa cta att act act tct gat aaa gaa			144
	Trp Leu Lys Asp Gln Gly His Glu Leu Ile Thr Thr Ser Asp Lys Glu			
	35 40 45			
	ggt gaa aca agt gaa ttg gat aaa cat atc cca gat gct gat att atc			192
	Gly Glu Thr Ser Glu Leu Asp Lys His Ile Pro Asp Ala Asp Ile Ile			
	50 55 60			

	atc acc act cct ttc cat cct gct tat atc act aag gaa aga ctt gac	240
	Ile Thr Thr Pro Phe His Pro Ala Tyr Ile Thr Lys Glu Arg Leu Asp	
	65 70 75 80	
5	aag gct aag aac tta aaa tta gtc gtt gtc gct ggt gtt ggt tct gat	288
	Lys Ala Lys Asn Leu Lys Leu Val Val Val Ala Gly Val Gly Ser Asp	
	85 90 95	
10	cac att gat tta gat tat att aat caa aca ggt aag aaa atc tca gtc	336
	His Ile Asp Leu Asp Tyr Ile Asn Gln Thr Gly Lys Lys Ile Ser Val	
	100 105 110	
15	ctg gaa gtt aca ggt tct aat gtt gtc tct gtt gct gaa cac gtt gtc	384
	Leu Glu Val Thr Gly Ser Asn Val Val Ser Val Ala Glu His Val Val	
	115 120 125	
20	atg acc atg ctt gtc ttg gtt aga aat ttc gtt cca gca cat gaa caa	432
	Met Thr Met Leu Val Leu Val Arg Asn Phe Val Pro Ala His Glu Gln	
	130 135 140	
25	att att aac cac gat tgg gag gtt gct gct atc gct aag gat gct tac	480
	Ile Ile Asn His Asp Trp Glu Val Ala Ala Ile Ala Lys Asp Ala Tyr	
	145 150 155 160	
30	gat atc gaa ggt aaa act atc gct acc att ggt gct ggt aga att ggt	528
	Asp Ile Glu Gly Lys Thr Ile Ala Thr Ile Gly Ala Gly Arg Ile Gly	
	165 170 175	
35	tac aga gtc ttg gaa aga tta ctc cca ttt aat cca aaa gaa tta tta	576
	Tyr Arg Val Leu Glu Arg Leu Leu Pro Phe Asn Pro Lys Glu Leu Leu	
	180 185 190	
40	tac tac gat tat caa gct tta cca aaa gaa gct gaa gaa aaa gtt ggt	624
	Tyr Tyr Asp Tyr Gln Ala Leu Pro Lys Glu Ala Glu Glu Lys Val Gly	
	195 200 205	
45	gct aga aga gtt gaa aat att gaa gaa tta gtt gct caa gct gat atc	672
	Ala Arg Arg Val Glu Asn Ile Glu Glu Leu Val Ala Gln Ala Asp Ile	
	210 215 220	
50	ggt aca gtt aat gct cca tta cac gca ggt aca aaa ggt tta att aat	720
	Val Thr Val Asn Ala Pro Leu His Ala Gly Thr Lys Gly Leu Ile Asn	
	225 230 235 240	
55	aag gaa tta tta tct aaa ttt aaa aaa ggt gct tgg tta gtc aat acc	768
	Lys Glu Leu Leu Ser Lys Phe Lys Lys Gly Ala Trp Leu Val Asn Thr	
	245 250 255	
60	gca aga ggt gct att gct gtt gct gaa gat gtt gca gca gct tta gaa	816
	Ala Arg Gly Ala Ile Ala Val Ala Glu Asp Val Ala Ala Ala Leu Glu	
	260 265 270	
65	tct ggt caa tta aga ggt tac ggt ggt gat gtt tgg ttc cca caa cca	864
	Ser Gly Gln Leu Arg Gly Tyr Gly Gly Asp Val Trp Phe Pro Gln Pro	
	275 280 285	
70	gct cca aag gat cac cca tgg aga gat atg aga aat aaa tat ggt gct	912
	Ala Pro Lys Asp His Pro Trp Arg Asp Met Arg Asn Lys Tyr Gly Ala	
	290 295 300	

	ggt aat gcc atg act cct cac tac tct ggt act act tta gac gct caa	960
	Gly Asn Ala Met Thr Pro His Tyr Ser Gly Thr Thr Leu Asp Ala Gln	
5	305 310 315 320	
	aca aga tac gct gaa ggt act aaa aat att ttg gaa tca ttc ttt acc	1008
	Thr Arg Tyr Ala Glu Gly Thr Lys Asn Ile Leu Glu Ser Phe Phe Thr	
	325 330 335	
10	ggt aaa ttt gat tac aga cca caa gat att atc tta tta aat ggt gaa	1056
	Gly Lys Phe Asp Tyr Arg Pro Gln Asp Ile Ile Leu Leu Asn Gly Glu	
	340 345 350	
15	tac gtt act aaa gct tac ggt aaa cac gat aag aaa taa	1095
	Tyr Val Thr Lys Ala Tyr Gly Lys His Asp Lys Lys	
	355 360	
20	<210> 8	
	<211> 364	
	<212> PRT	
	<213> Candida boidinii	
25	<400> 8	
	Met Lys Ile Val Leu Val Leu Tyr Asp Ala Gly Lys His Ala Ala Asp	
	1 5 10 15	
30	Glu Glu Lys Leu Tyr Gly Ser Thr Glu Asn Lys Leu Gly Ile Ala Asn	
	20 25 30	
35	Trp Leu Lys Asp Gln Gly His Glu Leu Ile Thr Thr Ser Asp Lys Glu	
	35 40 45	
40	Gly Glu Thr Ser Glu Leu Asp Lys His Ile Pro Asp Ala Asp Ile Ile	
	50 55 60	
45	Ile Thr Thr Pro Phe His Pro Ala Tyr Ile Thr Lys Glu Arg Leu Asp	
	65 70 75 80	
50	Lys Ala Lys Asn Leu Lys Leu Val Val Val Ala Gly Val Gly Ser Asp	
	85 90 95	
55	His Ile Asp Leu Asp Tyr Ile Asn Gln Thr Gly Lys Lys Ile Ser Val	
	100 105 110	
	Leu Glu Val Thr Gly Ser Asn Val Val Ser Val Ala Glu His Val Val	
	115 120 125	
	Met Thr Met Leu Val Leu Val Arg Asn Phe Val Pro Ala His Glu Gln	
	130 135 140	

5 Ile Ile Asn His Asp Trp Glu Val Ala Ala Ile Ala Lys Asp Ala Tyr  
 145 150 155 160  
 Asp Ile Glu Gly Lys Thr Ile Ala Thr Ile Gly Ala Gly Arg Ile Gly  
 165 170 175  
 10 Tyr Arg Val Leu Glu Arg Leu Leu Pro Phe Asn Pro Lys Glu Leu Leu  
 180 185 190  
 15 Tyr Tyr Asp Tyr Gln Ala Leu Pro Lys Glu Ala Glu Glu Lys Val Gly  
 195 200 205  
 20 Ala Arg Arg Val Glu Asn Ile Glu Glu Leu Val Ala Gln Ala Asp Ile  
 210 215 220  
 25 Val Thr Val Asn Ala Pro Leu His Ala Gly Thr Lys Gly Leu Ile Asn  
 225 230 235 240  
 Lys Glu Leu Leu Ser Lys Phe Lys Lys Gly Ala Trp Leu Val Asn Thr  
 245 250 255  
 30 Ala Arg Gly Ala Ile Ala Val Ala Glu Asp Val Ala Ala Ala Leu Glu  
 260 265 270  
 35 Ser Gly Gln Leu Arg Gly Tyr Gly Gly Asp Val Trp Phe Pro Gln Pro  
 275 280 285  
 40 Ala Pro Lys Asp His Pro Trp Arg Asp Met Arg Asn Lys Tyr Gly Ala  
 290 295 300  
 45 Gly Asn Ala Met Thr Pro His Tyr Ser Gly Thr Thr Leu Asp Ala Gln  
 305 310 315 320  
 Thr Arg Tyr Ala Glu Gly Thr Lys Asn Ile Leu Glu Ser Phe Phe Thr  
 325 330 335  
 50 Gly Lys Phe Asp Tyr Arg Pro Gln Asp Ile Ile Leu Leu Asn Gly Glu  
 340 345 350  
 55 Tyr Val Thr Lys Ala Tyr Gly Lys His Asp Lys Lys  
 355 360

&lt;211&gt; 5686

&lt;212&gt; DNA

&lt;213&gt; Artificial

5 &lt;220&gt;

&lt;223&gt; Plasmid pAM3.25

&lt;400&gt; 9

10	tatgaagatt gtcttagttc tttatgatgc tggtaagcac gctgctgatg aagaaaaatt	60
	atatggttct actgaaaata aattaggtat tgctaattgg ttaaaagatc aaggcatga	120
	actaattact acttctgata aagaaggatga aacaagtga ttggataaac atatcccaga	180
15	tgctgatatt atcatcacca ctcttttcca tcttgcttat atcactaagg aaagacttga	240
	caaggctaag aacttaaaat tagtcgttgt cgctgggtgtt ggttctgatc acattgattt	300
20	agattatatt aatcaaacag gtaagaaaat ctcatgctg gaagttacag gttctaattgt	360
	tgtctctgtt gctgaacacg ttgtcatgac catgcttgct ttggttagaa atttcgttcc	420
	agcacatgaa caaattatta accacgattg ggagggtgct gctatcgcta aggatgctta	480
25	cgatatcgaa ggtaaaaacta tcgctaccat tgggtgctggt agaattgggtt acagagtctt	540
	ggaaagatta ctcccattta atccaaaaga attattatac tacgattatc aagctttacc	600
30	aaaagaagct gaagaaaaag ttggtgctag aagagttgaa aatattgaag aattagttgc	660
	tcaagctgat atcgttacag ttaatgctcc attacacgca ggtacaaaag gtttaattaa	720
	taaggaatta ttatctaaat ttaaaaaagg tgcttggtta gtcaataccg caagagggtgc	780
35	tattgctggt gctgaagatg ttgcagcagc tttagaatct ggtcaattaa gaggttacgg	840
	tggtgatggt ttggtccac aaccagctcc aaaggatcac ccatggagag atatgagaaa	900
40	taaatatggt gctggtaatg ccatgactcc tcactactct ggtactactt tagacgctca	960
	aacaagatac gctgaaggta ctaaaaatat tttggaatca ttctttaccg gtaaatgtga	1020
	ttacagacca caagatatta tcttattaaa tggatgaatac gttactaaag cttacggtaa	1080
45	acacgataag aaataagacg tcaagcttgg ctgttttggc ggatgagaga agattttcag	1140
	cctgatacag attaaatcag aacgcagaag cggctctgata aaacagaatt tgcttgccgg	1200
50	cagtagcgcg gtggtccac ctgaccccat gccgaactca gaagtgaac gccgtagcgc	1260
	cgatggtagt gtggggtctc cccatgagag agtagggaac tgccaggcat caaataaaac	1320
	gaaaggctca gtcgaaagac tgggcctttc gttttatctg ttgtttgtcg gtgaacgctc	1380
55	tcctgagtag gacaaatccg ccgggagcgg atttgaacgt tgcgaagcaa cgccccggag	1440
	ggtggcgggc aggacgcccg ccataaactg ccaggcatca aattaagcag aaggccatcc	1500
	tgacggatgg cctttttgcg tttctacaaa ctcttttgtt tatttttcta aatacattca	1560

	aatatgtatc cgctcatgag acaataaccc tgataaatgc ttcaataata ttgaaaaagg	1620
5	aagagtatga gtattcaaca tttccgtgtc gcccttattc ctttttttgc ggcattttgc	1680
	cttcctgttt ttgctcacc agaaacgctg gtgaaagtaa aagatgctga agatcagttg	1740
	ggtgcacgag tgggttacat cgaactggat ctcaacagcg gtaagatcct tgagagtttt	1800
10	cgccccgaag aacgttttcc aatgatgagc actttttaaag ttctgctatg tggcgcggta	1860
	ttatcccgtg ttgacgccgg gcaagagcaa ctcggtcgcc gcatacacta ttctcagaat	1920
15	gacttggttg agtactcacc agtcacagaa aagcatctta cggatggcat gacagtaaga	1980
	gaattatgca gtgctgccat aaccatgagt gataacactg cggccaactt acttctgaca	2040
	acgatcggag gaccgaagga gctaaccgct tttttgcaca acatggggga tcatgtaact	2100
20	cgccttgatc gttgggaacc ggagctgaat gaagccatac caaacgacga gcgtgacacc	2160
	acgatgcctg tagcaatggc aacaacgttg cgcaaactat taactggcga actacttact	2220
25	ctagcttccc ggcaacaatt aatagactgg atggaggcgg ataaagttgc aggaccactt	2280
	ctgcgctcgg cccttccggc tggctggttt attgctgata aatctggagc cggtgagcgt	2340
	gggtctcgcg gtatcattgc agcactgggg ccagatggta agccctcccg tatcgtagtt	2400
30	atctacacga cggggagtcg ggcaactatg gatgaacgaa atagacagat cgctgagata	2460
	ggtgcctcac tgattaagca ttggtaactg tcagaccaag ttactcata tatactttag	2520
35	attgatttaa aacttcattt ttaattttaa aggatctagg tgaagatcct ttttgataat	2580
	ctcatgacca aaatccctta acgtgagttt tcgttccact gagcgtcaga ccccgtagaa	2640
	aagatcaaag gatcttcttg agatcctttt tttctgcgcg taatctgctg cttgcaaaca	2700
40	aaaaaaccac cgctaccagc ggtggtttgt ttgccggatc aagagctacc aactcttttt	2760
	ccgaaggtaa ctggcttcag cagagcgcag ataccaaata ctgtccttct agtgtagccg	2820
45	tagttaggcc accacttcaa gaactctgta gcaccgccta catacctcgc tctgctaate	2880
	ctgttaccag tggctgctgc cagtggcgat aagtcgtgtc ttaccggggt ggactcaaga	2940
	cgatagttac cggataaggc gcagcggtcg ggctgaacgg ggggttcgtg cacacagccc	3000
50	agcttggagc gaacgaccta caccgaactg agatacctac agcgtgagct atgagaaagc	3060
	gccacgcttc ccgaagggag aaaggcggac aggtatccgg taagcggcag ggtcggaaca	3120
55	ggagagcgca cgagggagct tccaggggga aacgcctggt atctttatag tctgtcggg	3180
	tttcgccacc tctgacttga gcgtcgattt ttgtgatgct cgtcaggggg gcggagccta	3240
	tggaaaaacg ccagcaacgc ggccttttta cggttcctgg ccttttgctg gccttttgct	3300



	cacatgttct ttcctgcgtt atcccctgat tctgtggata accgtattac cgcctttgag	3360
	tgagctgata ccgctcgccg cagccgaacg accgagcgca gcgagtcagt gagcgaggaa	3420
5	gcggaagagc gcctgatgcg gtattttctc cttacgcac tgtgcggtat ttcacaccgc	3480
	atatatggtg cactctcagt acaatctgct ctgatgccg atagttaagc cagtatacac	3540
10	tccgctatcg ctacgtgact gggtcacggc tgcgccccga caccgcgcaa caccgctga	3600
	cgcgcctga cgggcttgtc tgctccccgc atccgcttac agacaagctg tgaccgtctc	3660
	cgggagctgc atgtgtcaga ggttttcacc gtcacaccg aaacgcgcga ggcagctgcg	3720
15	gtaaagctca tcagcgtggc cgtgaagcga ttcacagatg tctgcctggt catccgcgtc	3780
	cagctcgttg agtttctcca gaagcgtaa tgtctggctt ctgataaagc gggccatggt	3840
20	aagggcggtt ttttctggt tggtcacttg atgcctccgt gtaaggggga atttctgttc	3900
	atgggggtaa tgataccgat gaaacgagag aggatgctca cgatacgggt tactgatgat	3960
	gaacatgcc gggtactgga acgttgtgag ggtaaacaac tggcggtatg gatgcggcgg	4020
25	gaccagagaa aaatcactca gggccaatgc cagcgcttcg ttaatacaga tgtagggtgtt	4080
	ccacagggta gccagcagca tcctgcgatg cagatccgga acataatggt gcagggcgct	4140
30	gacttcgcg tttccagact ttacgaaaca cggaaaccga agaccattca tgttggtgct	4200
	caggtcgcag acgttttgca gcagcagtcg cttcacgttc gctcgcgtat cggtgattca	4260
	ttctgctaac cagtaaggca accccgccag cctagccggg tcctcaacga caggagcacg	4320
35	atcatgcgca cccgtggcca ggacccaacg ctgcccga tgcgccgct gcggctgctg	4380
	gagatggcgg acgcgatgga tatgttctgc caagggttg tttgcgcatt cacagttctc	4440
40	cgcaagaatt gattggctcc aattcttggga gtggtgaatc cgtagcgag gtgccgccg	4500
	cttcattca ggtcgaggtg gccgggctcc atgcaccgc acgcaacgcg gggaggcaga	4560
	caaggtatag ggcggcgctt acaatccatg ccaaccggtt ccatgtgctc gccgaggcgg	4620
45	cataaatcgc cgtgacgatc agcgggtccag tgatcgaagt taggctggta agagccgcga	4680
	gcgatccttg aagctgtccc tgatggctcg catctacctg cctggacagc atggcctgca	4740
50	acgcgggcat cccgatgccg ccggaagcga gaagaatcat aatggggaag gccatccagc	4800
	ctcgcgtgc gaacgccagc aagacgtagc ccagcgcgtc ggccgccatg ccggcgataa	4860
	tggcctgctt ctcgccgaaa cgtttggtg cgggaccagt gacgaaggct tgagcgaggg	4920
55	cgtgcaagat tccgaatacc gcaagcgaca ggccgatcat cgtcgcgtc cagcgaaagc	4980
	ggtcctcgcc gaaaatgacc cagagcgctg ccggcacctg tcctacgagt tgcataataa	5040
	agaagacagt cataagtgcg gcgacgatag tcatgccccg cgccaccgg aaggagctga	5100

ctggggttgaa ggctctcaag ggcacgcgtc gacgctctcc cttatgacgac tccctgcatta 5160  
 5 ggaagcagcc cagtagtagg ttgaggccgt tgagcaccgc cgccgcaagg aatgggtgcat 5220  
 gctcgatggc tacgagggca gacagtaagt ggatttacca taatccctta attgtacgca 5280  
 ccgctaaaac gcgttcagcg cgatcacggc agcagacagg taaaaatggc aacaaaccac 5340  
 10 cctaaaaact gcgcgatcgc gcctgataaa ttttaaccgt atgaatacct atgcaaccag 5400  
 aggggtacagg ccacattacc cccacttaat ccactgaagc tgccattttt catgggtttca 5460  
 15 ccatcccagc gaagggccat gcatgcacgc aaattaatac gacgaaatta atacgactca 5520  
 ctatagggca attgcatca ccacaattca gcaaattgtg aacatcatca cgttcatctt 5580  
 tccctgggtg ccaatggccc attttctgt cagtaacgag aaggctcgca attcaggcgc 5640  
 20 tttttagact ggctgtaatg aacaattctt aagaaggaga tataca 5686  
  
 <210> 10  
 <211> 5106  
 25 <212> DNA  
 <213> Artificial  
  
 <220>  
 <223> Plasmid pAM10.1  
 30  
 <400> 10  
 gaaggagata tacatatgac attagaaatc ttgaataact tagaaaaata tgattatgag 60  
 35 caagtagtat tttgtcaaga taaagaatct ggtttaaaag caattattgc aattcatgat 120  
 acaacacttg gaccggctct tgggtgaaca agaatgtgga catatgattc tgaagaagcg 180  
 gcgattgaag atgcattgcg tcttgcaaaa gggatgacat aaaaaaacgc agcagctggc 240  
 40 ttaaacttag gtggtgcgaa aacagtaatt atcggtgatc ctcgtaaaga taagagcgaa 300  
 gcaatgttcc gtgcactagg acgttatatc caaggactaa acggacgtta cattacagct 360  
 45 gaagatgttg gtacaacagt agatgatatg gatattatcc atgaagaaac tgactttgta 420  
 acaggtatct caccatcatt cggttcttct ggtaacccat ctccggtaac tgcatacggc 480  
 gtttaccgtg gtatgaaagc agctgcaaaa gaagctttcg gtactgacaa tttagaagga 540  
 50 aaagtaattg ctgttcaagg cgttggtaac gtagcatatc acctatgcaa acatttacac 600  
 gctgaaggag caaaattaat tgttacagat attaataaag aagctgtaca acgtgctgta 660  
 55 gaagaattcg gtgcacagc agttgaacca aatgaaattt acgggtgttg atgcgatatt 720  
 tacgcaccat gtgcactagg cgcaacagtt aatgatgaaa ctattccaca acttaaagca 780  
 aaagtaatcg caggttctgc gaataaccaa ttaaaagaag atcgatcatg tgacatcatt 840

	catgaaatgg gtattgtata cgcaccagat tatgtaatta atgcaggtgg cgtaattaac	900
	gtagcagacg aattatatgg atacaataga gaacgtgcac taaaacgtgt tgagtctatt	960
5	tatgacacga ttgcaaaagt aatcgaaatt tcaaaacgcg atggcatagc aacttatgta	1020
	gcggcagatc gtctagctga agagcgcatt gcaagcttga agaattctcg tagcacttac	1080
10	ttacgcaacg gtcacgatat tattagccgt cgctaacgcg tttgcggttg gcaaaatggc	1140
	gcagcagcaa ggcgtggcgg tgaaaacctc tgccgaagcc ctgcaacagg ccattgacga	1200
	taatttcttg caagccgaat accgcgacta ccgccgtacc tccatctaaa agcttatcga	1260
15	tgataagctg tcaaacatga gaattacaac ttatatcgta tggggctgac ttcaggtgct	1320
	acatttgaag agataaattg cactgaaatc tagaaatatt ttatctgatt aataagatga	1380
20	tcttcttgag atcgtttttg tctgcgcgta atctcttgct ctgaaaacga aaaaaccgcc	1440
	ttgcagggcg gtttttcgaa ggttctctga gctaccaact ctttgaaccg aggtaactgg	1500
	cttgaggag cgcagtcacc aaaacttgtc ctttcagttt agccttaacc ggcgcatgac	1560
25	ttcaagacta actcctctaa atcaattacc agtggctgct gccagtgggtg cttttgcatg	1620
	tctttccggg ttggactcaa gacgatagtt accggataag gcgcagcggc cggactgaac	1680
30	gggggggttcg tgcatacagt ccagcttgga gcgaactgcc tacccggaac tgagtgtcag	1740
	gcgtggaatg agacaaacgc ggccataaca gcggaatgac accggtaaac cgaaaggcag	1800
	gaacaggaga gcgcacgagg gagccgccag gggaaacgcc tggatatctt atagtcctgt	1860
35	cgggtttcgc caccactgat ttgagcgtca gatttcgtga tgcttgctcag gggggcggag	1920
	cctatggaaa aacggctttg ccgcggccct ctacttccc tgttaagtat ctccctggca	1980
40	tcttccagga aatctccgcc ccgttcgtaa gccatttccg ctgcgccgag tcgaacgacc	2040
	gagcgtagcg agtcagttag cgaggaagcg gaatatatcc tgtatcacat attctgctga	2100
	cgcaccggtg cagccttttt tctcctgcca catgaagcac ttcactgaca ccctcatcag	2160
45	tgccaacata gtaagccagt atacactccg ctagcgtga tgtccggcgg tgcttttgcc	2220
	gttacgcacc accccgtcag tagctgaaca ggaggacag ctgatagaaa cagaagccac	2280
50	tggagcacct caaaaacacc atcatacact aaatcagtaa gttggcagca tcaccgcagc	2340
	cactttgcgc cgaataaata cctgtgacgg aagatcactt cgcagaataa ataaatcctg	2400
	gtgtccctgt tgataccggg aagccctggg ccaacttttg gcgaaaatga gacgttgatc	2460
55	ggcacgtaag aggttccaac tttcaccata atgaaataag atcactaccg ggcgtatttt	2520
	ttgagttatc gagattttca ggagctaagg aagctaaaat ggagaaaaaa atcactggat	2580
	ataccaccgt tgatatatcc caatggcatc gtaaagaaca ttttgaggca tttcagtcag	2640

	ttgctcaatg tacctataac cagaccgttc agctggatat tacggccttt ttaaagaccg	2700
5	taaagaaaaa taagcacaag ttttatccgg cctttattca cattcttgcc cgctgatga	2760
	atgctcatcc ggaattccgt atggcaatga aagacgggtga gctgggtgata tgggatagtg	2820
	ttcacccttg ttacaccggt ttccatgagc aaactgaaac gttttcatcg ctctggagtg	2880
10	aataccacga cgatttccgg cagtttctac acatatattc gcaagatgtg gcgtgttacg	2940
	gtgaaaacct ggcctatttc cctaaagggt ttattgagaa tatgtttttc gtctcagcca	3000
15	atccctgggt gagtttcacc agttttgatt taaacgtggc caatatggac aacttcttcg	3060
	ccccgtttt caccatgggc aaatattata cgcaaggcga caagggtgctg atgccgctgg	3120
	cgattcaggt tcatcatgcc gtctgtgatg gcttccatgt cggcagaatg cttaatgaat	3180
20	tacaacagta ctgcgatgag tggcagggcg gggcgtaatt tttttaaggc agttattggt	3240
	gcccttaaac gcctggtgct acgcctgaat aagtgataat aagcggatga atggcagaaa	3300
25	ttcgaaagca aattcgaccc ggtcgtcgggt tcagggcagg gtcgttaaat agccgcttat	3360
	gtctattgct ggtttaccgg tttattgact accggaagca gtgtgaccgt gtgctttctca	3420
	aatgcctgag gccagtttgc tcaggctctc cccgtggagg taataattga cgatatgac	3480
30	atttattctg cctcccagag cctgataaaa acggttagcg cttcgttaat acagatgtag	3540
	gtgttcaca gggtagccag cagcatcctg cgatgcagat ccggaacata atgggtgcagg	3600
35	gcgcttgttt cggcgtgggt atgggtggcag gccccgtggc cgggggactg ttgggcgctg	3660
	ccggcacctg tcctacgagt tgcattgataa agaagacagt cataagtgcg gcgacgatag	3720
	tcattgccccg cgcccaccgg aaggagctac cggacagcgg tgcggactgt tgtaactcag	3780
40	aataagaaat gagggcgctc atggcggttg ctctcagtca tagtatcgtg gtatcacccg	3840
	ttggttccac tctctgttgc gggcaacttc agcagcacgt aggggacttc cgcgtttcca	3900
45	gactttacga aacacggaaa ccgaagacca ttcatgttgt tgctcaggtc gcagacgttt	3960
	tgcagcagca gtcgcttcac gttcgctcgc gtatcgggtga ttcattctgc taaccagtaa	4020
	ggcaaccccc ccagcctagc cgggtcctca acgacaggag cacgatcatg cgcacccgtg	4080
50	gccaggaccc aacgctgccc gagatgcgcc gcgtgcggct gctggagatg gcggacgcga	4140
	tggatatgtt ctgccaaggg ttggtttgcg cattcacagt tctccgcaag aattgattgg	4200
55	ctccaattct tggagtgggt aatccgttag cgaggtgccg ccggcttcca ttcaggtcga	4260
	ggtggccccg ctccatgcac cgcgacgcaa cgcggggagg cagacaagggt atagggcggc	4320
	gcctacaatc catgccaacc cgttccatgt gctcgccgag gcggcataaa tcgccgtgac	4380

gatcagcggg ccagtgatcg aagttaggct ggtaagagcc gcgagcgatc cttgaagctg 4440  
 tccctgatgg tcgtcatcta cctgcctgga cagcatggcc tgcaacgcgg gcatcccgat 4500  
 5 gccgcccggaa gcgagaagaa tcataatggg gaaggccatc cagcctcgcg tcgcgaacgc 4560  
 cagcaagacg tagcccagcg cgtcggccgc catgccggcg ataatggcct gcttctcgcc 4620  
 10 gaaacgtttg gtggcgggac cagtgcgaa ggcttgagcg agggcggtgca agattccgaa 4680  
 taccgcaagc gacaggccga tcatcgtcgc gctccagcga aagcggtcct cgccgaaaat 4740  
 gacccagagc gctgccggca cctgtcctac gagttgcatg ataaagaaga cagtcataag 4800  
 15 tgcgggcgacg atagtcatgc cccgcgcca ccggaaggag ctgactgggt tgaaggctct 4860  
 caagggcatc ggtcgacgct ctcccttatg cgactcctgc attaggaagc agcccagtag 4920  
 20 taggttgagg ccgttgagca ccgcccgcgc aaggaatggg gcatgcatcg atcaccacaa 4980  
 ttcagcaa at tgtgaacatc atcacgttca tctttccctg gttgccaatg gccattttc 5040  
 ctgtcagtaa cgagaaggc gcgaattcag gcgcttttta gactggtcgt aatgaacaat 5100  
 25 tcttaa 5106

<210> 11  
 <211> 5597  
 30 <212> DNA  
 <213> Unknown

<220>  
 <223> Plasmid  
 35

<220>  
 <221> CDS  
 <222> (25)..(1749)  
 <223> scfA - malic enzyme gene  
 40

<400> 11  
 aattcttaag aaggagatat acat atg gat att caa aaa aga gtg agt gac 51  
 45 Met Asp Ile Gln Lys Arg Val Ser Asp  
 1 5

atg gaa cca aaa aca aaa aaa cag cgt tcg ctt tat atc cct tac gct 99  
 Met Glu Pro Lys Thr Lys Lys Gln Arg Ser Leu Tyr Ile Pro Tyr Ala  
 10 15 20 25

50 ggc cct gta ctg ctg gaa ttt ccg ttg ttg aat aaa ggc agt gcc ttc 147  
 Gly Pro Val Leu Leu Glu Phe Pro Leu Leu Asn Lys Gly Ser Ala Phe  
 30 35 40

55 agc atg gaa gaa cgc cgt aac ttc aac ctg ctg ggg tta ctg ccg gaa 195  
 Ser Met Glu Glu Arg Arg Asn Phe Asn Leu Leu Gly Leu Leu Pro Glu  
 45 50 55

gtg gtc gaa acc atc gaa gaa caa gcg gaa cga gca tgg atc cag tat 243

	Val Val Glu Thr Ile Glu Glu Gln Ala Glu Arg Ala Trp Ile Gln Tyr	
	60 65 70	
5	cag gga ttc aaa acc gaa atc gac aaa cac atc tac ctg cgt aac atc Gln Gly Phe Lys Thr Glu Ile Asp Lys His Ile Tyr Leu Arg Asn Ile	291
	75 80 85	
10	cag gac act aac gaa acc ctc ttc tac cgt ctg gta aac aat cat ctt Gln Asp Thr Asn Glu Thr Leu Phe Tyr Arg Leu Val Asn Asn His Leu	339
	90 95 100 105	
15	gat gag atg atg cct gtt att tat acc cca acc gtc ggc gca gcc tgt Asp Glu Met Met Pro Val Ile Tyr Thr Pro Thr Val Gly Ala Ala Cys	387
	110 115 120	
	gag cgt ttt tct gag atc tac cgc cgt tca cgc ggc gtg ttt atc tct Glu Arg Phe Ser Glu Ile Tyr Arg Arg Ser Arg Gly Val Phe Ile Ser	435
	125 130 135	
20	tac cag aac cgg cac aat atg gac gat att ctg caa aac gtg ccg aac Tyr Gln Asn Arg His Asn Met Asp Asp Ile Leu Gln Asn Val Pro Asn	483
	140 145 150	
25	cat aat att aaa gtg att gtg gtg act gac ggt gaa cgc att ctg ggg His Asn Ile Lys Val Ile Val Val Thr Asp Gly Glu Arg Ile Leu Gly	531
	155 160 165	
30	ctt ggt gac cag ggc atc ggc ggg atg ggc att ccg atc ggt aaa ctg Leu Gly Asp Gln Gly Ile Gly Gly Met Gly Ile Pro Ile Gly Lys Leu	579
	170 175 180 185	
35	tcg ctc tat acc gcc tgt ggc ggc atc agc ccg gcg tat acc ctt ccg Ser Leu Tyr Thr Ala Cys Gly Gly Ile Ser Pro Ala Tyr Thr Leu Pro	627
	190 195 200	
	gtg gtg ctg gat gtc gga acg aac aac caa cag ctg ctt aac gat ccg Val Val Leu Asp Val Gly Thr Asn Asn Gln Gln Leu Leu Asn Asp Pro	675
	205 210 215	
40	ctg tat atg ggc tgg cgt aat ccg cgt atc act gac gac gaa tac tat Leu Tyr Met Gly Trp Arg Asn Pro Arg Ile Thr Asp Asp Glu Tyr Tyr	723
	220 225 230	
45	gaa ttc gtt gat gaa ttt atc cag gct gtg aaa caa cgc tgg cca gac Glu Phe Val Asp Glu Phe Ile Gln Ala Val Lys Gln Arg Trp Pro Asp	771
	235 240 245	
50	gtg ctg ttg cag ttt gaa gac ttt gct caa aaa aat gcg atg ccg tta Val Leu Leu Gln Phe Glu Asp Phe Ala Gln Lys Asn Ala Met Pro Leu	819
	250 255 260 265	
55	ctt aac cgc tat cgc aat gaa att tgt tct ttt aac gat gac att cag Leu Asn Arg Tyr Arg Asn Glu Ile Cys Ser Phe Asn Asp Asp Ile Gln	867
	270 275 280	
	ggc act gcg gcg gta aca gtc ggc aca ctg atc gca gca agc cgc gcg Gly Thr Ala Ala Val Thr Val Gly Thr Leu Ile Ala Ala Ser Arg Ala	915
	285 290 295	

	gca ggt ggt cag tta agc gag aaa aaa atc gtc ttc ctt ggc gca ggt Ala Gly Gln Leu Ser Glu Lys Lys Ile Val Phe Leu Gly Ala Gly 300 305 310	963
5	tca gcg gga tgc ggc att gcc gaa atg atc atc tcc cag acc cag cgc Ser Ala Gly Cys Gly Ile Ala Glu Met Ile Ile Ser Gln Thr Gln Arg 315 320 325	1011
10	gaa gga tta agc gag gaa gcg gcg cgg cag aaa gtc ttt atg gtc gat Glu Gly Leu Ser Glu Glu Ala Ala Arg Gln Lys Val Phe Met Val Asp 330 335 340 345	1059
15	cgc ttt ggc ttg ctg act gac aag atg ccg aac ctg ctg cct ttc cag Arg Phe Gly Leu Leu Thr Asp Lys Met Pro Asn Leu Leu Pro Phe Gln 350 355 360	1107
20	acc aaa ctg gtg cag aag cgc gaa aac ctc agt gac tgg gat acc gac Thr Lys Leu Val Gln Lys Arg Glu Asn Leu Ser Asp Trp Asp Thr Asp 365 370 375	1155
	agc gat gtg ctg tca ctg ctg gat gtg gtg cgc aat gta aaa cca gat Ser Asp Val Leu Ser Leu Leu Asp Val Val Arg Asn Val Lys Pro Asp 380 385 390	1203
25	att ctg att ggc gtc tca gga cag acc ggg ctg ttt acg gaa gag atc Ile Leu Ile Gly Val Ser Gly Gln Thr Gly Leu Phe Thr Glu Glu Ile 395 400 405	1251
30	atc cgt gag atg cat aaa cac tgt ccg cgt ccg atc gtg atg ccg ctg Ile Arg Glu Met His Lys His Cys Pro Arg Pro Ile Val Met Pro Leu 410 415 420 425	1299
35	tct aac ccg acg tca cgc gtg gaa gcc aca ccg cag gac att atc gcc Ser Asn Pro Thr Ser Arg Val Glu Ala Thr Pro Gln Asp Ile Ile Ala 430 435 440	1347
40	tgg acc gaa ggt aac gcg ctg gtc gcc acg ggc agc ccg ttt aat cca Trp Thr Glu Gly Asn Ala Leu Val Ala Thr Gly Ser Pro Phe Asn Pro 445 450 455	1395
	gtg gta tgg aaa gat aaa atc tac cct atc gcc cag tgt aac aac gcc Val Val Trp Lys Asp Lys Ile Tyr Pro Ile Ala Gln Cys Asn Asn Ala 460 465 470	1443
45	ttt att ttc ccg ggc atc ggc ctg ggt gtt att gct tcc ggc gcg tca Phe Ile Phe Pro Gly Ile Gly Leu Gly Val Ile Ala Ser Gly Ala Ser 475 480 485	1491
50	cgt atc acc gat gag atg ctg atg tgc gca agt gaa acg ctg gcg cag Arg Ile Thr Asp Glu Met Leu Met Ser Ala Ser Glu Thr Leu Ala Gln 490 495 500 505	1539
55	tat tca cca ttg gtg ctg aac ggc gaa ggt atg gta ctg ccg gaa ctg Tyr Ser Pro Leu Val Leu Asn Gly Glu Gly Met Val Leu Pro Glu Leu 510 515 520	1587
	aaa gat att cag aaa gtc tcc cgc gca att gcg ttt gcg gtt ggc aaa Lys Asp Ile Gln Lys Val Ser Arg Ala Ile Ala Phe Ala Val Gly Lys 525 530 535	1635

	atg gcg cag cag caa ggc gtg gcg gtg aaa acc tct gcc gaa gcc ctg	1683
	Met Ala Gln Gln Gln Gly Val Ala Val Lys Thr Ser Ala Glu Ala Leu	
	540 545 550	
5	caa cag gcc att gac gat aat ttc tgg caa gcc gaa tac cgc gac tac	1731
	Gln Gln Ala Ile Asp Asp Asn Phe Trp Gln Ala Glu Tyr Arg Asp Tyr	
	555 560 565	
10	cgc cgt acc tcc atc taa aagcttatcg atgataagct gtcaaacatg	1779
	Arg Arg Thr Ser Ile	
	570	
15	agaattacaa cttatatcgt atggggctga cttcaggtgc tacatttgaa gagataaatt	1839
	gcactgaaat ctagaaatat tttatctgat taataagatg atcttcttga gatcgttttg	1899
	gtctgcgcgt aatctcttgc tctgaaaacg aaaaaaccgc cttgcagggc ggtttttcga	1959
20	aggttctctg agctaccaac tctttgaacc gaggttaactg gcttggagga gcgcagtcac	2019
	caaaacttgt cctttcagtt tagccttaac cggcgcacatga cttcaagact aactcctcta	2079
25	aatcaattac cagtggctgc tgccagtggg gcttttgcac gtctttccgg gttggactca	2139
	agacgatagt taccggataa ggcgacagcg tccgactgaa cgggggggttc gtgcatacag	2199
	tccagcttgg agcgaactgc ctaccggaa ctgagtgtca ggcgtggaat gagacaaacg	2259
30	cggccataac agcggaatga caccggtaaa ccgaaaggca ggaacaggag agcgcacgag	2319
	ggagccgcca ggggaaacgc ctggtatctt tatagtcctg tcgggtttcg ccaccactga	2379
35	tttgagcgtc agatttcgtg atgcttgtca ggggggcgga gcctatggaa aaacggcttt	2439
	gccgcggccc tctcacttcc ctgttaagta tcttcttggc atcttccagg aaatctccgc	2499
	cccgttcgta agccatttcc gctcgccgca gtcgaacgac cgagcgtagc gagtcagtga	2559
40	gcgaggaagc ggaatatatc ctgtatcaca tattctgctg acgcaccggt gcagcctttt	2619
	ttctctgcc acatgaagca cttcactgac accctcatca gtgccaacat agtaagccag	2679
45	tatacactcc gctagcgtg atgtccggcg gtgcttttgc cgttacgcac cacccttca	2739
	gtagctgaac aggagggaca gctgatagaa acagaagcca ctggagcacc tcaaaaacac	2799
	catcatacac taaatcagta agttggcagc atcaccgcac gcactttgcg ccgaataaat	2859
50	acctgtgacg gaagatcact tcgcagaata aataaatcct ggtgtccctg ttgataccgg	2919
	gaagccctgg gccaaactttt ggcgaaaatg agacgttgat cggcacgtaa gaggttccaa	2979
55	ctttcaccat aatgaaataa gatcactacc gggcgtattt ttgagttat cgagattttc	3039
	aggagctaag gaagctaaaa tggagaaaaa aatcactgga tataccaccg ttgatataac	3099
	ccaatggcat cgtaaagaac attttgaggc atttcagtca gttgctcaat gtacctataa	3159



ccagaccggt cagctggata ttacggcctt tttaaagacc gtaaagaaaa ataagcacia 3219  
 gttttatccg gcctttattc acattcttgc ccgcctgatg aatgctcacc cggaattccg 3279  
 5 tatggcaatg aaagacgggtg agctgggtgat atgggatagt gttcaccctt gttacaccgt 3339  
 tttccatgag caaactgaaa cgtttttcacc gctctggagt gaataccacc acgattttccg 3399  
 10 gcagttttcta cacatatatt cgcaagatgt ggctgtgttac ggtgaaaacc tggcctatatt 3459  
 ccctaaaggg tttattgaga atatgttttt cgtctcagcc aatccctggg tgagttttcac 3519  
 cagttttgat ttaaactgtg ccaatatgga caacttcttc gcccccggtt tcaccatggg 3579  
 15 caaatattat acgcaaggcg acaagggtgct gatgccgctg gcgattcagg ttcacatgc 3639  
 cgtctgtgat ggcttccatg tcggcagaat gcttaatgaa ttacaacagt actgcgatga 3699  
 20 gtggcagggc ggggcgtaat ttttttaagg cagttattgg tgcccttaaa cgcttggtgc 3759  
 tacgcctgaa taagtataa taagcggatg aatggcagaa attcgaaagc aaattcgacc 3819  
 cggctgctcg ttcagggcag ggtcggttaa tagccgctta tgtctattgc tggtttaccg 3879  
 25 gttttattgac taccggaagc agtgtgaccg tgtgcttctc aaatgcctga ggccagtttg 3939  
 ctcaggctct ccccggtggag gtaataattg acgatatgat catttattct gcctcccaga 3999  
 30 gcctgataaa aacgggttagc gcttcgttaa tacagatgta ggtgttccac agggtagcca 4059  
 gcagcatcct gcgatgcaga tccggaacat aatgggtgcag ggcgcttggt tcggcgtggg 4119  
 tatgggtggca gggcccggtg ccgggggact gttgggcgtt gccggcacct gtcctacgag 4179  
 35 ttgcatgata aagaagacag tcataagtgc ggcgacgata gtcatgcccc gcgcccaccg 4239  
 gaaggagcta ccggacagcg gtgcggactg ttgtaactca gaataagaaa tgaggccgct 4299  
 40 catggcggtt actctcagtc atagtatcgt ggtatcaccg gttggttcca ctctctgttg 4359  
 cgggcaactt cagcagcagc taggggactt ccgcgtttcc agactttacc aaacacggaa 4419  
 accgaagacc attcatgttg ttgctcaggt ccgacagcgt ttgcagcagc agtcgcttca 4479  
 45 cgttcgctcg cgtatcggtg attcattctg ctaaccagta aggcaacccc gccagcctag 4539  
 ccgggtcctc aacgacagga gcacgatcat gcgcaccggt ggccaggacc caacgctgcc 4599  
 50 cgagatgcgc cgctgctggc tgctggagat ggcgacgcg atggatatgt tctgccaagg 4659  
 gttggtttgc gcattcacag ttctccgcaa gaattgattg gctccaattc ttggagtggg 4719  
 gaatccgtta gcgaggtgcc gccggcttcc attcaggtcg aggtggcccc gctccatgca 4779  
 55 ccgcgacgca acgcggggag gcagacaagg tatagggcgg cgcttacaat ccatgccaac 4839  
 ccgttccatg tgctcgccga ggcggcataa atcgccgtga cgatcagcgg tccagtgate 4899  
 gaagttaggc tggtaagagc ccgagcgcat ccttgaagct gtccctgatg gtcgtcatct 4959

acctgcctgg acagcatggc ctgcaacgcg ggcattccga tgccgccgga agcgagaaga 5019  
 5 atcataatgg ggaaggccat ccagcctcgc gtcgcgaacg ccagcaagac gtagcccagc 5079  
 gcgtcggccg ccatgccggc gataatggcc tgcttctcgc cgaaacgttt ggtggcggga 5139  
 ccagtgaaga aggcttgagc gagggcgtgc aagattccga ataccgcaag cgacaggccg 5199  
 10 atcatcgtcg cgctccagcg aaagcgggcc tcgccgaaaa tgaccagag cgctgccggc 5259  
 acctgtccta cgagttgcat gataaagaag acagtcataa gtgcggcgac gatagtcatg 5319  
 15 ccccgcgccc accggaagga gctgactggg ttgaaggctc tcaagggcat cggtcgacgc 5379  
 tctcccttat gcgactcctg cattaggaag cagcccagta gtaggttgag gccgttgagc 5439  
 accgccgccc caaggaatgg tgcattgcat gatcaccaca attcagcaaa ttgtgaacat 5499  
 20 catcacgttc atctttccct ggttgccaat ggccatttt cctgtcagta acgagaaggt 5559  
 cgccaattca ggcgcttttt agactggctg taatgaac 5597  
  
 25 <210> 12  
 <211> 574  
 <212> PRT  
 <213> Unknown  
  
 30 <220>  
 <223> Plasmid  
  
 <400> 12  
  
 35 Met Asp Ile Gln Lys Arg Val Ser Asp Met Glu Pro Lys Thr Lys Lys  
 1 5 10 15  
  
 40 Gln Arg Ser Leu Tyr Ile Pro Tyr Ala Gly Pro Val Leu Leu Glu Phe  
 20 25 30  
  
 Pro Leu Leu Asn Lys Gly Ser Ala Phe Ser Met Glu Glu Arg Arg Asn  
 35 40 45  
 45  
  
 Phe Asn Leu Leu Gly Leu Leu Pro Glu Val Val Glu Thr Ile Glu Glu  
 50 55 60  
  
 50 Gln Ala Glu Arg Ala Trp Ile Gln Tyr Gln Gly Phe Lys Thr Glu Ile  
 65 70 75 80  
  
 55 Asp Lys His Ile Tyr Leu Arg Asn Ile Gln Asp Thr Asn Glu Thr Leu  
 85 90 95  
  
 Phe Tyr Arg Leu Val Asn Asn His Leu Asp Glu Met Met Pro Val Ile

100

105

110

5 Tyr Thr Pro Thr Val Gly Ala Ala Cys Glu Arg Phe Ser Glu Ile Tyr  
115 120 125

10 Arg Arg Ser Arg Gly Val Phe Ile Ser Tyr Gln Asn Arg His Asn Met  
130 135 140

15 Asp Asp Ile Leu Gln Asn Val Pro Asn His Asn Ile Lys Val Ile Val  
145 150 155 160

Val Thr Asp Gly Glu Arg Ile Leu Gly Leu Gly Asp Gln Gly Ile Gly  
165 170 175

20 Gly Met Gly Ile Pro Ile Gly Lys Leu Ser Leu Tyr Thr Ala Cys Gly  
180 185 190

25 Gly Ile Ser Pro Ala Tyr Thr Leu Pro Val Val Leu Asp Val Gly Thr  
195 200 205

30 Asn Asn Gln Gln Leu Leu Asn Asp Pro Leu Tyr Met Gly Trp Arg Asn  
210 215 220

Pro Arg Ile Thr Asp Asp Glu Tyr Tyr Glu Phe Val Asp Glu Phe Ile  
225 230 235 240

35 Gln Ala Val Lys Gln Arg Trp Pro Asp Val Leu Leu Gln Phe Glu Asp  
245 250 255

40 Phe Ala Gln Lys Asn Ala Met Pro Leu Leu Asn Arg Tyr Arg Asn Glu  
260 265 270

45 Ile Cys Ser Phe Asn Asp Asp Ile Gln Gly Thr Ala Ala Val Thr Val  
275 280 285

50 Gly Thr Leu Ile Ala Ala Ser Arg Ala Ala Gly Gly Gln Leu Ser Glu  
290 295 300

Lys Lys Ile Val Phe Leu Gly Ala Gly Ser Ala Gly Cys Gly Ile Ala  
305 310 315 320

55 Glu Met Ile Ile Ser Gln Thr Gln Arg Glu Gly Leu Ser Glu Glu Ala  
325 330 335

Ala Arg Gln Lys Val Phe Met Val Asp Arg Phe Gly Leu Leu Thr Asp  
340 345 350

5 Lys Met Pro Asn Leu Leu Pro Phe Gln Thr Lys Leu Val Gln Lys Arg  
355 360 365

10 Glu Asn Leu Ser Asp Trp Asp Thr Asp Ser Asp Val Leu Ser Leu Leu  
370 375 380

15 Asp Val Val Arg Asn Val Lys Pro Asp Ile Leu Ile Gly Val Ser Gly  
385 390 395 400

20 Gln Thr Gly Leu Phe Thr Glu Glu Ile Ile Arg Glu Met His Lys His  
405 410 415

Cys Pro Arg Pro Ile Val Met Pro Leu Ser Asn Pro Thr Ser Arg Val  
420 425 430

25 Glu Ala Thr Pro Gln Asp Ile Ile Ala Trp Thr Glu Gly Asn Ala Leu  
435 440 445

30 Val Ala Thr Gly Ser Pro Phe Asn Pro Val Val Trp Lys Asp Lys Ile  
450 455 460

35 Tyr Pro Ile Ala Gln Cys Asn Asn Ala Phe Ile Phe Pro Gly Ile Gly  
465 470 475 480

40 Leu Gly Val Ile Ala Ser Gly Ala Ser Arg Ile Thr Asp Glu Met Leu  
485 490 495

Met Ser Ala Ser Glu Thr Leu Ala Gln Tyr Ser Pro Leu Val Leu Asn  
500 505 510

45 Gly Glu Gly Met Val Leu Pro Glu Leu Lys Asp Ile Gln Lys Val Ser  
515 520 525

50 Arg Ala Ile Ala Phe Ala Val Gly Lys Met Ala Gln Gln Gln Gly Val  
530 535 540

55 Ala Val Lys Thr Ser Ala Glu Ala Leu Gln Gln Ala Ile Asp Asp Asn  
545 550 555 560

Phe Trp Gln Ala Glu Tyr Arg Asp Tyr Arg Arg Thr Ser Ile  
565 570